

# Triadimefon 500

Unique to 4Farmers

<b>Active constituent</b>	Triadimefon 500g/kg	<b>Formulation</b> WP	
<b>Typical situations</b>	Wheat and barle.		
<b>Diseases controlled</b>	Barley Powdery Mildew, Wheat Leaf and Stripe Rust and Take-all (suppression).		
<b>Chemical group</b>	3		
<b>Mode of action</b>	Upwardly systemic in plant—provides protectant, curative and eradivative activity.		
<b>Typical pack size</b>	20kg		
<b>Poison schedule</b> 6	<b>Dangerous goods class</b> –	<b>UN</b> –	<b>Packing code</b> –
<b>Withholding period</b>	<b>Harvest</b> Do not harvest for 3 months post application.		
	<b>Grazing</b> Do not graze or cut treated foliage for stock food.		
<b>Application method</b>	Mixed dry with fertiliser and applied in-furrow.		<b>Rain fastness</b> N/A
<b>Efficacy</b>	Apply product evenly onto the fertiliser. Ideally the powder should be metered via an applicator into an auger or elevator.		
<b>Adjuvants</b>	None Required		
<b>Compatibility</b>			
<b>Incompatibility</b>			
<b>Water quality</b>			
<b>4F Broadacre registrations</b>	Wheat and barley		
<b>4F Other registrations</b>	N/A.		
<b>Similar product registrations</b>	N/A.		
<b>Situation</b>	<b>Target diseases</b>	<b>Rate/ha</b>	<b>Comments</b>
Barley	Powdery Mildew	200g	Use when disease risk is high, and monitor crops from 10-12 weeks for follow up control requirements. Up to 16 weeks control not uncommon.
Wheat	Leaf rust, stripe rust		
	<b>Suppression:</b> Take All Speckled leaf blotch		
<b>Comments</b>			
<ul style="list-style-type: none"> <li>Used as a fertiliser treatment, Triadimefon provides cost effective protectant activity for the above diseases; when residual activity declines and/or where disease persists, another product should be used for curative activity.</li> <li><b>Care:</b> May impact coleoptile length in some circumstances; ensure adequate seed/fertiliser separation, particularly when sown into pre-emergent herbicide treated soil.</li> <li>Dangerous to aquatic organisms and soil fungi (temporary).</li> </ul>			