

Linuron 450

Similar to: Linurex®

Active constituent	Linuron 450g/L	Formulation SC	
Typical situations	Wheat, barley, oats and other (see registration section).		
Chemical group	5 (C)		
Mode of action	Inhibits photosynthesis.		
Typical pack size	10L		
Poison schedule –	Dangerous goods class –	UN –	Packing code –
Withholding period	Harvest Cereals: Withholding period not required when used as directed. Horticulture: Various, see label.		
Plant back	–		
Application method	Boom spray 80L–110L/ ha and aerial 20L–30L/ ha.	Rain fastness 24 hrs	
Efficacy	Do not apply to weeds under severe stress.		
Adjuvants	Do not add adjuvants.		
Compatibility	4Farmers Diclofop-Methyl.		
Incompatibility	Wetting agents, surfactants, and crop oils.		
Water quality	–		
Time to effects and symptoms	5 days desiccation begins.		
4F Broadacre registrations	Wheat, barley, oats.		
4F Other registrations	Various horticultural crops, coriander seed crops.		
Similar product registrations	–		
Situation	Target Weed/s	Rate/ha	Comments
Wheat, barley, oats (Z13–Z14)	Doublegee, capeweed, wild radish, corn gromwell, wild turnip, deadnettle, mustard, turnip weed, rough poppy, slender thistle	600 ml	All states.
		300 ml + (MCPA 500g/L) 400 ml	WA only. Crops generally 3 to 4-leaf (Z13–Z14).
	Amsinckia, saffron thistle	950 ml	All states.
Comments			
<ul style="list-style-type: none"> Do not allow mixture to stand in vat for prolonged periods, as it is difficult to re-suspend and may lose effectiveness. Do not add 4Farmers Diclofop-Methyl or any herbicide to this product + MCPA. Do not apply with misting machines. 			
			