

Butafenacil 200 EC

Similar to: *B in Logran B®*

Active constituent	Butafenacil 200g/L	Formulation EC	
Typical situations	For the control of marshmallow in fallow and prior to establishment of cereal crops.		
Chemical group	14 (G)		
Mode of action	Inhibitor of protoporphyrinogen oxidase.		
Typical pack size	5L		
Poison schedule –	Dangerous goods class –	UN –	Packing code III
Withholding period	Harvest Not required when used as directed.		
	Grazing Not required when used as directed.		
	Fallow Do not graze or cut sprayed vegetation for stock food for at least 1 day after application.		
	Cereals Do not graze or cut for stock food for 6 weeks after application.		
Plant back	N/A		
Application method	Boomspray only. Do not apply by aircraft.		Rain fastness 2 hrs
Efficacy	Ensure thorough coverage. Do not apply if runoff is expected within 3 days. Do not sow or cultivate for 1 hour after spraying. Don't use disk sowing systems or ones that allow treated soil to contact seed of broad-leaved crops.		
Adjuvants	Wetting agents: Usually improve performance. Spray oils: Usually improve performance.		
Compatibility	Paraquat + Diquat, 360g/L Paraquat, 250g/L Paraquat, Prosulfocarb, Trifluralin, Triallate, Triasulfuron, Atrazine, Simazine, S-metolachlor, Metribuzin, Diuron, Isoxaflutole, 2,4-D Ester 680.		
Incompatibility	4Farmers Butafenacil must not be tank mixed with other formulations of Butafenacil.		
Water quality	–		
Time to effects and symptoms	–		
4F Broadacre registrations	Fallow.		
4F Other registrations	–		
Similar product registrations	–		

Situation	Target Weed/s	Rate/ha	Comments
Fallow	Marshmallow	60–80ml/ha + Glyphosate at labelled rates	Do not make more than 1 application per season. Rate Selection: Use the higher rate where weeds are stressed, there is significant overlap of target weed leaves, where targeting larger weeds or when applying in Spring or Summer. Always use an adjuvant at recommended label rates. Esterified vegetable oil (e.g. 4Farmers Speedy Spray), crop oil concentrate (e.g. 4Farmers Take Up), or a non-ionic wetter (e.g. 4Farmers Wetter 1000)

Comments

- **Do not** irrigate to the point of run-off for at least 48 hours after application.
- **Do not** spray weeds which are waterlogged, under stress of any kind or covered with soil or dust.
- **Do not** spray weeds covered with heavy dew.
- **Do not** use prior to the sowing of crops undersown with pasture species or vetch.
- **Do not** tank mix with other formulations of Butafenacil.
- **Do not** use if other formulations of Butafenacil are to be used in the same season.

