Basher® Similar to: Slasher®

Active constituent	Nonanoic or pelargonic acid 525 g/L		Formulation EC	
Typical situations	Organic pre-sowing knockdown herbicide.			
Chemical group	0			
Mode of action	Knockdown.			
Typical pack size	1L, 5L 20L, 110L, 1000L			
Poison schedule -	Dangerous goods class -	UN -	Packing code -	
Withholding period	Harvest Not required when used as directed.			
	Grazing As above, except when tank mixed with compatible products (see label).			
Plant back	None.			
Application method	Boom or spot spray.		Rain fastness 24hours	
Efficacy	Half fill spray tank and commence agitation. Slowly add required amount of Basher herbicide.			
Adjuvants	0.1% wetter.			
Compatibility	Often used in organic situations and so additions with other chemicals is uncommon.			
Incompatibility	N/A—as above			
Water quality	Limited issues.			
Time to effects and symptoms	Discolouration occurs within 4 days.			
4F Broadacre registrat	ions -			
4F Other registrations	-			

Situation	Target Weed/s	Rate/ha	Comments
Outdoor areas— Paths, driveways, gardens, near sheds, amenity areas, protected cropping situations, near nursery and orchids. Spot spraying in lawns	Seedling and young annual and perennial weeds and grasses including fat hen, potato weed, stinking roger, couch grass, chickweed, lamb's tongue, peppercress, capeweed, common storksbill, clover.  More advanced or established weeds may only be suppressed and require repeated application.	70ml per litre of water	Spray to completely cover weed surface. Reapplication at around 7 days may be necessary if regrowth occurs. This product may damage lawn and turf when used as a spot spray.  Perennial weed species may require repeated applications to obtain long term control.
and turf. Fallows.  General use— Including near buildings, commercial and industrial areas, domestic and public service areas, rights-of way, bushland and forests.		Mix 7L per 100L water Apply 900– 1400 L/ha	To control weeds in Protected cropping situations, apply on each side of the crop row taking care not to allow spray to contact any of the crop foliage.
Gardens, paths, rockeries, tiled.	Algae, lichen, liverwort, moss	70ml per litre of water	Spray to fully cover plant surface. For driveways or heavy infestations, clean off by brushing after 3 days and reapply if necessary.

## Comments

· Wear eye protection when mixing or using.

Similar product registrations Slasher®

- · Do not enter treated areas until the product has dried.
- Avoid drift and do not contaminate water ways.

