

# Dicyclanil 50

Similar to: CLiK®

Active constituent	Dicyclanil 50g/L	Formulation SC	
Chemical group	–		
Mode of action	–		
Typical pack size	20L		
Poison schedule –	Dangerous goods class –	UN –	Packing code –
Application method	Use only a spray-on applicator gun.		Rain fastness –
Usage	For the protection of sheep against fly strike from blow fly ( <i>Lucilia cuprina</i> ) for 18–24 weeks. Also, for the protection of mulesing and marking wounds on sheep against fly strike during healing process.		

Situation	Animal weight	Rate	Comments
Poll strike prevention	–	5ml	Apply to the areas requiring protection, and around poll and horns.
Body strike prevention	10–30kgs	7–8ml x 2 bands	Ensure bands overlap up and down the midline.
	31–50+ kgs	10–12ml x 2 bands	
Breech strike prevention	10–30kgs	7–8ml	–
	31–50+ kgs	10–12ml x 1 band	
After mulesing prevention	Up to 30kgs	4ml x 2 bands	Apply two bands, one on each side of the open wound with small overlap onto surrounding wool.
	31–50+ kgs	5ml x 2 bands	
After marking prevention	Up to 30kgs (ram lambs)	4ml x 2 bands	When the tail has been amputated, apply 4ml over the crutch to ensure sufficient coverage over the wound. When the tail has not been amputated, ensure sufficient coverage of the tail where rubber rings is applied.
	Up to 30kgs (ewe lambs)	4ml x 1 band	

## Comments

- Read label before use.
- This product is not recommended for treating existing strikes.
- Heavy rain following application of this product could diminish the period of protection.
- **Withholding periods:**
  - Do not use less than 28 days before slaughter for human consumption
  - Do not use on sheep which are producing or may in the future produce milk or milk products for human consumption
  - Do not use less than 3 months before shearing.

