

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



## BUTROXYDIM 250 WG

Herbicide

ACTIVE CONSTITUENT: 250 g/L BUTROXYDIM

GROUP A HERBICIDE

For the control of certain grasses in a range of broadacre crops as per the Directions for Use.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

4 FARMERS AUSTRALIA PTY LTD

A.B.N. 51 160 092 428

1 / 70 McDowell Street, Welshpool, WA 6106

Tel: (08) 9356 3445 Fax: (08) 9356 3447

Batch Number: \_\_\_\_\_ Date of Manufacture: \_\_\_\_\_  
APVMA Approval No: 85134/112344

### DIRECTIONS FOR USE

#### RESTRAINTS

DO NOT treat weeds that are not actively growing or are growing under stress. Under such circumstances the biological processes of the weeds slow down and 4Farmers Butroxydim 250 WG transport in the weed can be drastically reduced, resulting in an incomplete kill or suppression only of the weeds. DO NOT allow annual grass weeds to reach the late tillering stage before application.

DO NOT apply at flowering stage of weeds.

DO NOT apply under conditions of prolonged high or very low temperature (frosts), moisture stress (waterlogging or drought), low humidity, poor fertility or stress from previous herbicide application.

Crop	Weeds Controlled	Weed Stage	State	Rate/Ha	Critical Comments
Chickpeas Faba beans Field peas Lentils Linseed Lucerne Lupins Vetches ( <i>Vicia</i> spp.) White and Subterranean Clover and Medic pastures	Annual Ryegrass ( <i>Lolium rigidum</i> ) Barley Grass ( <i>Hordeum</i> spp.) Wild Oats ( <i>Avena</i> spp.)  Annual Ryegrass ( <i>Lolium rigidum</i> ) together with any of the following grasses: Annual Phalaris ( <i>Phalaris</i> spp.) Barley Grass ( <i>Hordeum</i> spp.) Brome Grass ( <i>Bromus</i> spp.) Volunteer Cereals – Wheat ( <i>Triticum aestivum</i> ) – Barley ( <i>Hordeum vulgare</i> ) – Oats ( <i>Avena sativa</i> ) – Triticale ( <i>x Triticosecale</i> ) Wild Oats ( <i>Avena</i> spp.)	2 leaf to early tillering Zadoks 12-23	All States	80 to 180g *  80 to 180g *  # plus an effective rate of a fop herbicide containing fluazifop, haloxyfop, propaquizafop, or quizalofop	Use the lower rates for younger weeds growing actively under ideal conditions. Use the higher rates for weeds that are predominantly at early tillering, or where denser populations are present or under less than ideal growing conditions or where some weed resistance to Group A herbicides is suspected. Because of the extreme variation of cross resistance present in annual ryegrass populations there is no guarantee that 4Farmers Butroxydim 250 WG will always provide consistently high levels of control but trials have shown that higher rates may be expected to perform better than lower rates. For all crops apply in not less than 50L water/ha.  # Refer to the label of the partner herbicide for rates and specific directions for use. 4Farmers Butroxydim 250 WG has good activity on barley grass and wild oats but is weaker on Brome Grass and Volunteer Cereals so the addition of a partner fop herbicide is generally recommended where any of these other weeds occur together with Annual Ryegrass. For the partner herbicide use rates at the lowest end of the range recommended for the particular grass weed. Guidance can also be obtained from labelled tank mixes of the partner herbicide with other dim herbicides.
Canola (conventional, TT, "IMI" varieties)	Annual Ryegrass ( <i>Lolium rigidum</i> ) together with any of the following grasses Annual Phalaris ( <i>Phalaris</i> spp.) Barley Grass ( <i>Hordeum</i> spp.) Brome Grass ( <i>Bromus</i> spp.) Volunteer Cereals – Wheat ( <i>Triticum aestivum</i> ) – Barley ( <i>Hordeum vulgare</i> ) – Oats ( <i>Avena sativa</i> ) – Triticale ( <i>x Triticosecale</i> ) Wild oats ( <i>Avena</i> spp.)	2 leaf to early tillering Zadoks 12-23	All States	80g*  # plus an effective rate of a fop herbicide containing fluazifop, haloxyfop, propaquizafop, quizalofop	<b>Warning</b> Canola can be sensitive to 4Farmers Butroxydim 250 WG. DO NOT use more than 80g/ha on canola. To minimize leaf symptoms do not apply to any variety before the majority of plants (80%) are at the true 4 leaf stage (4th leaf expanded, 5th leaf emerging). Under certain conditions, all varieties may exhibit leaf symptoms and/or reduced early competitiveness. If this occurs canola yield is usually unaffected but in some instances may be reduced. Avoid spraying stressed crops as crops growing under stress will show slower recovery. Avoid spraying practices that lead to over application such as double overlap, and spraying out corners. # Refer to section above for control of additional grasses.
Lucerne Mung beans Navy beans Peanuts Soybeans	<b>Echinochloa</b> species – Awnless Barnyard Grass ( <i>E. colona</i> ) – Barnyard Grass ( <i>E. crus-galli</i> ) – Japanese Millet ( <i>E. utilis</i> ) <b>Digitaria</b> species – Crab Grass ( <i>D. sanguinalis</i> ) – Summer Grass ( <i>D. ciliaris</i> ) <b>Chloris</b> species – Feather Top Rhodes Grass ( <i>C. virgata</i> ) – Windmill Grass ( <i>C. truncate</i> ) <b>Setaria</b> species – Dwarf Setaria ( <i>S. italica</i> ) – Whorled Pigeon Grass ( <i>S. verticillate</i> )	2 leaf to early tillering Zadoks 12-23	Qld, NSW, Vic & NT only	120g or 180g *	Use the lower rate for the control of seedling grasses at the pre-tillering growth stage and growing under good conditions. Use the higher rate for control of grasses at the early tillering (2 to 3 tillers) growth stage. Aerial application – see Spraying Instructions.
Cotton	<b>Brachiaria</b> species – Green Summer Grass ( <i>B. subquadripara</i> ) – Velvet Grass ( <i>B. windersii</i> ) <b>Volunteer Crops</b> – Maize ( <i>Zea mays</i> ) – Sorghum ( <i>Sorghum bicolor</i> ) Crowfoot Grass ( <i>Eleusine indica</i> ) Dinebra ( <i>Dinebra reflexa</i> ) Early Spring Grass ( <i>Eriochloa pseudoacrotricha</i> ) Johnson Grass ( <i>Sorghum halepense</i> ) (seedling)		Qld & NSW only		
Sunflowers	Liverseed Grass ( <i>Urochloa panicoides</i> ) Spiny Burr Grass ( <i>Cenchrus incertus</i> )  <b>Eragrostis</b> species – Elastic Grass ( <i>E. tenuifolia</i> ) – Mexican Love Grass ( <i>E. mexicana</i> ) – Stink Grass ( <i>E. cilianensis</i> )  <b>Volunteer cereals</b> – Wheat ( <i>Triticum aestivum</i> ) – Barley ( <i>Hordeum vulgare</i> )  <b>Other Grasses</b> Coast Button Grass ( <i>Dactyloctenium aegyptium</i> ) Grader Grass ( <i>Themeda quadrivalvis</i> )	2 leaf to 5 leaf but prior to tillering  2 leaf to early tillering Zadoks 12-22, prior to stem elongation  2 leaf to 5 leaf but prior to tillering	Qld, NSW & Vic only	120g *  180g *  180g *	For the control of pre-tillering grasses only. Do not apply to tillered grasses.  For the control of grasses from 2 leaf to early tillering only (max. 2 tillers), prior to stem elongation or booting.  Use for suppression only of pre-tillered grasses. Do not apply to tillered weeds.

\* Always apply with 4Farmers Turbo Charge at 1L/100L of spray solution. For aerial application apply 4Farmers Turbo Charge at a rate of 1L/ha.

Note: Does NOT control winter grass (*Poa annua*), Silver grass (*Vulpia* spp.), Nutsedge (*Cyperus* spp.) and broadleaf weeds.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

#### WITHHOLDING PERIODS

WITHHOLDING PERIOD PRIOR TO HARVEST NOT REQUIRED WHEN USED AS DIRECTED

Canola, Chick peas, Clover, Faba beans, Field peas, Lentils, Linseed, Lucerne, Lupins, Medics, Mung beans, Navy beans, Peanuts, Soybeans, Sunflowers, Vetches:  
DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

Cotton: DO NOT GRAZE OR CUT FOR STOCKFEED. DO NOT FEED COTTON TRASH TO LIVESTOCK.

## GENERAL INSTRUCTIONS

### Stress conditions

Moisture stress is a particular problem, whether caused by drought or short term stresses, resulting from a combination of high temperatures and low humidity or by prior use of pre-emergence herbicides resulting in stunted root growth of weeds. An irrigation or effective rainfall must occur immediately before or after spraying if these conditions exist. Other stress conditions to avoid include use of pre-emergence herbicides resulting in stunted root growth of weeds, frost, waterlogging and extended cold conditions. Do not spray if any stress conditions are evident just prior to application.

### Mixing

Add the required amount of 4Farmers Butroxydim 250 WG (see table) directly to the spray tank which should be 50% full with clean water. After the product is fully dispersed complete filling the tank. The agitation system should be running during mixing and spraying. If agitation ceases and settling occurs, re-suspend contents thoroughly before spraying. Ensure that all in-line strainer and nozzle screens in the sprayer are 100 mesh or coarser.

Add any other product next.

An oil adjuvant is essential and should be added last to the spray tank.

4Farmers Turbo Charge at 1 L/100L of water is the only recommended adjuvant for ground application. For aerial application use 4Farmers Turbo Charge at a rate of 1 L/ha.

4Farmers Turbo Charge must be added to at least 10 times its volume of water especially where a suction (filling) probe is used.

The area covered per measure pack will depend upon the application rate/ha according to the following table:

4Farmers Butroxydim 250 WG Rate/ha	Hectares					
	1 kg	2 kg	3 kg	4 kg	5 kg	10 kg
80g	12.5	25	37.5	50	62.5	125
100g	1	20	30	40	50	100
150g	6	13.3	20	26.7	33.3	66.7
180g	5	11.1	16.7	22.2	27.8	55.6

### Application

Time of application: 4Farmers Butroxydim 250 WG is rainfast within 30 minutes. Do not apply 4Farmers Butroxydim 250 WG if rain is expected within 30 minutes. To obtain the maximum benefits of this product, weeds should be sprayed when young and actively growing. Use the lower rates for grasses with up to 4 leaves. Use the higher rates for weeds with up to 3 tillers.

Unless otherwise specified in the Critical Comments application may be made at any stage of crop growth but allowing sufficient time for the prescribed withholding period.

**Method of application:** Good spray coverage is essential for maximum results. Spray equipment must be checked and calibrated accurately prior to application.

**Boom spraying:** Check height of boom above the target weeds to ensure the spray is evenly distributed and a double overlap pattern is obtained. An even distribution of droplets with a Fine spray quality (measured with water to ASAE S572 standard) is desirable for maximum efficacy. The inclusion of 4Farmers Turbo Charge coarsens the spray quality when applied through a TeeJet XR11002 nozzle such that a Medium spray quality results which will help reduce the likelihood of drift.

**Broadacre use:** Water volumes per hectare will depend on nozzle selection and ground speed but should be in the range of 50-100L/ha. 110° flat fan nozzles are preferred for use with this product. Use higher volumes of spray to achieve better coverage of dense, vigorous weed infestations and/or if the nozzles used produce coarser than a Fine spray quality (measured with water). The use of air induction nozzles is not recommended until further trial data is available.

**Aerial application:** The product may be applied through boom or Micronair units in 20-30L of water per hectare. The use of the higher volume is preferred due to more reliable results. Spray quality should be Fine (measured with water to ASAE S572 standard) and swath width should not be wider than recommended for type of plane and application unit. Weed control following aerial application may not be as good as from ground application.

### Sprayer cleanup or decontamination

Before spraying sensitive crops (which include canola, chickpeas, faba beans, field peas, lupins, lucerne, clover etc) thoroughly remove all traces of SU herbicides such as 4Farmers Metsulfuron 600, 4Farmers Chlorsulfuron 750 or 4Farmers Triasulfuron 750 or 4Farmers Carfentazone 240 from mixing and spray equipment immediately after use as recommended on the label for that herbicide.

### Sprayer cleanup or decontamination

Before spraying sensitive crops (which include canola, chickpeas, faba beans, field peas, lupins, lucerne, clover etc) thoroughly remove all traces of SU herbicides such as 4Farmers Metsulfuron 600, 4Farmers Chlorsulfuron 750 or 4Farmers Triasulfuron 750 or 4Farmers Carfentazone 240 from mixing and spray equipment immediately after use as recommended on the label for that herbicide.

### Compatibility

The following recommendations are based on results from field and laboratory tests with 4Farmers Butroxydim 250 WG plus 1% 4Farmers Turbo Charge. Compatibility is defined as acceptable physical mixing, nil or generally transient crop effect and grass control within 10% of that from 4Farmers Butroxydim 250 WG alone. Performance of any tank mix will be better if the maximum rate of 4Farmers Butroxydim 250 WG is used and the weed growth stages are according to the label. Always refer to the label of the partner product for rates and any specific guidelines and restraints for the crop or situation.

4Farmers Butroxydim 250 WG is compatible with any one of the following herbicides for the crops specified:

Canola – 4Farmers Atrazine 900WG, 4Farmers Simazine 900 WG, 4Farmers Clopyralid 300, 4Farmers Haloxypfop 520, 4Farmers Clethodim 240, 4Farmers Fluazifop, 4Farmers Quizalofop, propaquizafop.

Pasture, clover based (not brome grass) – 4Farmers Flumetsulam 800 WG, bromoxynil, 4Farmers LVE MCPA 570, (max rate 0.5 L/ha; some clover vigour reduction possible), 4Farmers LV MCPA/Diflufenican, 4Farmers Bromoxynil Diflufenican, 4Farmers Haloxypfop 520, 4Farmers Clethodim 240, 4Farmers Simazine 900 WG, 4Farmers Fluazifop, 4Farmers Quizalofop, propaquizafop.

4Farmers Butroxydim 250 WG is compatible with any one of the following insecticides for the crops specified:

Pulses, canola, pasture – Imidan, Karate, / Karate with Zeon / Matador with Zeon (lambda-cyhalothrin), 4Farmers Omethoate 290, 4Farmers Pirimicarb 500 WG, 4Farmers Dimethoate 400.

4Farmers Butroxydim 250 WG is compatible with any one of the following fungicides for the crops specified:

Pulses – 4Farmers Mancozeb 750, 4Farmers Procymidone 500.

4Farmers Butroxydim 250 WG is compatible with any one of the following trace elements for the crops specified:

Canola, Pulses, Pasture - EDTA chelates of Cu, Mn, Zn.

4Farmers Butroxydim 250 WG is not compatible with 4Farmers Chlorpyrifos 500 (reduced grass weed control); 4Farmers Diflufenican 500, or Eclipse (increased crop effect).

DO NOT tank mix 4Farmers Butroxydim 250 WG plus 4Farmers Turbo Charge with more than one of the above products before doing a jar test beforehand. However, physical compatibility does not guarantee biological compatibility. Do not tank mix with other products or trace elements without reference to a 4Farmers Australia Pty Ltd representative.

## RESISTANT WEEDS WARNING

### GROUP A HERBICIDE

4Farmers Butroxydim 250 WG Herbicide is a member of the Cyclohexanedione (DIM) group of herbicides. The product has the inhibitors of acetyl coA carboxylase mode of action. For weed resistance management, the product is a Group A herbicide.

Some naturally occurring weed biotypes resistant to the product and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group A herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, 4Farmers Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment which may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT plant cereal crops for a period of 4 weeks after application of 4Farmers Butroxydim 250 WG Herbicide.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Highly toxic to algae and aquatic plants. DO NOT contaminate dams, rivers or streams with the chemical or used containers.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Avoid inhaling dust. When preparing product for use wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## SAFETY DATA SHEET

For further information please refer to the Safety Data Sheet. For a copy visit our website at [www.4farmers.com.au](http://www.4farmers.com.au)

## CONDITIONS OF SALE

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. 4Farmers Australia Pty Ltd's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which 4Farmers Australia Pty Ltd shall have no liability.

<b>In a Transport Emergency Dial 000 Police or Fire Brigade</b>	<b>Special Poisons Advice: 13 11 26</b>
---	---