

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



TRIASULFURON 750 WG

Selective Herbicide

ACTIVE CONSTITUENT: 750 g/kg TRIASULFURON

GROUP B HERBICIDE

For pre-plant control of Annual Ryegrass, Paradoxa Grass and certain broadleaf weeds in Wheat, and for Post-Emergent control of Wild Radish in Wheat, Oats and Barley as per Directions for Use.

ALWAYS READ THE LABEL AND USE ONLY AS DIRECTED

4 FARMERS AUSTRALIA PTY LTD
A.B.N. 51 160 092 428
1 / 70 McDowell Street, Welshpool, WA 6106
Tel: (08) 9356 3445

Batch Number: _____ Date of Manufacture: _____

APVMA Approval No: 57753/48475

DIRECTIONS FOR USE

WESTERN AUSTRALIA ONLY – Pre-emergent application

RESTRAINTS

DO NOT apply to crops undersown with legumes.

DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Wheat	Burr Medic, Common Cotula, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Densiflower Fumitory, Hedge Mustard, Indian Hedge Mustard, Matricaria, Paterson's Curse, Rough Poppy, Smallflower Fumitory, Wards Weed, Wild Turnip, Yellow Burrweed (Amsinckia) and suppression of Crassula.	30g	Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10cm combine points. Application should not be made to ridged or excessively cloddy soil. For best results apply to moist soil when follow up rain is likely to occur within 7-10 days. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing. When used on Kulin wheat on very acid soils (pH 5.5 in water) or under poor fertility conditions increased stem breakage may occur. Early season crop retardation may occur where the product is used on soils with a pH greater than 8, and which are prone to zinc deficiency.
	Annual Ryegrass, Capeweed, Doublegee or Three-cornered Jack (Spiny Emex), Wireweed, and suppression of Wild Radish, and also 60-80% suppression of Soursob.	35g	
	Annual Ryegrass, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Wild Turnip, Wireweed (Hogweed), Ward's weed	10g to 15g plus 1 L of a 400g/L trifluralin	This mixture to be only used on alkaline soils where pH is greater than 8 (Water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and incorporate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall (greater than 50mm) within 7 days of application may affect efficacy especially at the 10g/ha rate. Late germination of some weeds eg. annual ryegrass will not be controlled in seasons of above average rainfall. Use the higher rate where heavy density of weeds is expected.

WESTERN AUSTRALIA ONLY – Post-emergent application

RESTRAINTS

DO NOT apply to crops undersown with legumes.

DO NOT spray when very dry conditions prevail.

DO NOT spray under dry frost conditions.

DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

IF RAIN FALLS within 6 hours of application, the effect could be diminished.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Barley, Oats, Wheat	Wild Radish	10g to 15g plus crop oil at 1L per 100L of spray mixture or surfactant at recommended label rates.	Early Post Emergent Application: Spray prior to the crop reaching mid-tillering (Zadoks – 23) and when wild radish is in the 2 to 6 leaf stage. Use rates towards the lower end of the range when weeds are small and growing conditions ideal. Spray only when weeds are actively growing. Late Post Emergent Application: Spray during early flowering. DO NOT apply to the crop during crop anthesis or flowering (Zadoks 60-69). DO NOT spray on weeds under stress. WARNING: Application to oats may cause some yellowing.
	Turnip Weed	6.5g + 300mL Terbutryn®, 500 SC	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for field peas, 10 leaflet for vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.
	Densiflower, Fumitory, Hedge Mustard, Smallflower Fumitory	6.5g + 300mL Terbutryn®, 500 SC to 10g + 440mL Terbutryn®, 500 SC	Spray only when weeds are actively growing.
	Ball Mustard, Field Pea (volunteer)	10g + 440mL Terbutryn®, 500 SC	
	Deadnettle	10g + 440mL Terbutryn®, 500 SC to 13g + 600mL Terbutryn®, 500 SC	Spray only after good rain and when top soil is moist. Best results are obtained when good soil moisture has been present since planting.
	Australian Crassula, Doublegee or Three-cornered Jack (Spiny Emex), Volunteer lupins	13g + 600mL Terbutryn®, 500 SC	

SOUTH AUSTRALIA ONLY – Pre-emergent application

RESTRAINTS

DO NOT apply to crops undersown with legumes.

DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Wheat	Ball Mustard, Burr Medic, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Densiflower Fumitory, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce, Rough Poppy, Smallflower Fumitory, Stemless Thistle, Vetch, Wards Weed, Wild Turnip, Yellow Burrweed (Amsinckia).	30g	Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10cm combine points. Application should not be made to ridged or excessively cloddy soil. For best results apply to moist soil when follow up rain is likely to occur within 7-10 days. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing. 4Farmers Triasulfuron will provide good control of volunteer grain legumes, however a small proportion of plants may survive and require an overspray to eliminate the potential for grain contamination. For Skeleton Weed a significant degree of control will be achieved on soil types of a predominantly sandy clay loam mixture with a pH greater than 8. Best control is observed where Skeleton Weed germinates in the very early stages of the crop. Surviving plants will be stunted.
	Annual Ryegrass, Capeweed, Three-cornered Jack or Doublegee (Spiny Emex), Volunteer Chickpeas, Faba Beans and Field Peas, Wireweed, suppression of Skeleton Weed, Wild Radish, 60-80% control of Soursob and suppression of remaining plants.	35g	
	Annual Ryegrass, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Wild Turnip, Wireweed (Hogweed), Ward's weed	10g to 15g plus 1 L of a 400g/L trifluralin	This mixture to be only used on alkaline soils where pH is greater than 8 (1:5 Soil:Water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and incorporate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall (greater than 50mm) within 7 days of application may affect efficacy especially at the 10g/ha rate. Late germination of some weeds eg. annual ryegrass will not be controlled in seasons of above average rainfall. Use the higher rate where heavy density of weeds is expected.

SOUTH AUSTRALIA ONLY – Post-emergent application

RESTRAINTS

DO NOT apply to crops undersown with legumes.

DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Barley, Oats, Wheat Late Post-emergent application	Wild Radish	15g plus crop oil at 1L per 100L of spray mixture or surfactant at recommended label rates.	Post Emergent Application: Spray during early flowering. DO NOT apply to the crop during crop anthesis or flowering (Zadoks 60-69). DO NOT spray on weeds under stress. WARNING: Application to oats may cause some yellowing.
	Faba Beans (volunteer), Long Fruited Turnip, Long Headed Poppy, Paterson's Curse, Turnip Weed	6.5g + 300mL Terbutryn®, 500 SC	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for field peas, 10 leaflet for vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.
	Hedge Mustard, Indian Hedge Mustard, Wild Turnip	6.5g + 300mL Terbutryn®, 500 SC to 10g + 440mL Terbutryn®, 500 SC	Spray only when weeds are actively growing.
	Ball Mustard, Crassula, Medic, Prickly Lettuce (Whipthistle)	10g + 440mL Terbutryn®, 500 SC	
Barley, Oats, Triticale, Wheat – from 3 leaf to early tillering stage	Field Pea (volunteer), Wild Radish	10g + 440mL Terbutryn®, 500 SC to 13g + 600mL Terbutryn®, 500 SC	Spray only after good rain and when top soil is moist. Best results are obtained when good soil moisture has been present since planting.
	Lupins (volunteer), Three-cornered Jack or Doublegee (Spiny Emex), Vetch, Wireweed (less than 3 leaves) suppression.	13g + 600mL Terbutryn®, 500 SC	

NEW SOUTH WALES & ACT ONLY – Pre-emergent application

RESTRAINTS

DO NOT apply to crops undersown with legumes.

DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Wheat	Black Bindweed (Climbing Buckwheat), Burr Medic, Common Cotula, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Densiflower Fumitory, Hedge Mustard, Indian Hedge Mustard, Lesser Swinecress, Paterson's Curse, Prickly Lettuce, Rough Poppy, Shepherd's Purse, Smallflower Fumitory, Sowthistle, Stagger Weed, Turnip Weed, Wild Turnip, Yellow Burrweed (Amsinckia) and suppression of Variegated Thistle.	30g	Apply to bare moist soil prior to sowing and incorporate by the sowing operation using low profile 10cm combine points. Application should not be made to ridged or excessively cloddy soil. For best results apply to moist soil when follow up rain is likely to occur within 7-10 days. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing.
	Annual Ryegrass, Capeweed, Doublegee or Three-cornered Jack (Spiny Emex), Paradoxa Grass (Annual Phalaris), Wireweed, and suppression of Wild Radish and Mexican Poppy.	35g	For best results for Paradoxa Grass control, apply to dry soil before the sowing rain.
	Annual Ryegrass, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Wild Turnip, Wireweed (Hogweed), Ward's weed	(Southern NSW only) 10g to 15g plus 1 L of a 400g/L trifluralin	This mixture to be only used on alkaline soils where pH is greater than 8 (1:5 Soil:Water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and incorporate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall (greater than 50mm) within 7 days of application may affect efficacy especially at the 10g/ha rate. Late germination of some weeds eg. annual ryegrass will not be controlled in seasons of above average rainfall. Use the higher rate where heavy density of weeds is expected.

NEW SOUTH WALES & ACT ONLY – Post-emergent application

RESTRAINTS

DO NOT apply to crops undersown with legumes.

DO NOT spray when very dry conditions prevail.

DO NOT spray under dry frost conditions.

DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

IF RAIN FALLS within 6 hours of application, the effect could be diminished.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Barley, Oats, Wheat Late Post-emergent application	Wild Radish	15g plus crop oil at 1L per 100L of spray mixture or surfactant at recommended label rates.	Late Post-emergent Application: Spray during early flowering. DO NOT apply to the crop during crop anthesis or flowering (Zadoks 60-69). DO NOT spray on weeds under stress. WARNING: Application to oats may cause some yellowing.
	Turnip Weed	6.5g + 300mL Terbutryn®, 500 SC	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for field peas, 10 leaflet for vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.
	Hedge Mustard, Indian Hedge Mustard, London Rocket, Wild Turnip	6.5g + 300mL Terbutryn®, 500 SC to 10g + 440mL Terbutryn®, 500 SC	Spray only when weeds are actively growing.
	Black Bindweed (Climbing Buckwheat), Faba Bean (volunteer), Field Pea (volunteer), Medic, Shepherds Purse	10g + 440mL Terbutryn®, 500 SC	Spray only after good rain and when top soil is moist.
	Deadnettle, Mexican Poppy, Wild Radish	10g + 440mL Terbutryn®, 500 SC to 13g + 600mL Terbutryn®, 500 SC	Best results are obtained when good soil moisture has been present since planting.
	Coreopsis, Sunflower (volunteer), Vetch	13g + 600mL Terbutryn®, 500 SC	

QUEENSLAND ONLY – Pre-emergent application

RESTRAINTS

DO NOT apply to crops undersown with legumes.

DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Wheat	African Turnip Weed, Black Bindweed (Climbing Buckwheat), Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, New Zealand Spinach, Prickly Lettuce, Slender Celery, Smallflower Fumitory, Turnip Weed, Yellow Burrweed (Amsinckia)	30g	Apply to weed free soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10cm combine points. Application should not be made to ridged or excessively cloddy soil. For best results apply when follow up rain is likely to occur within 7-10 days. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing. For Paradoxa Grass control, apply to dry soil before the sowing rain.
	Annual Ryegrass, Common Peppergrass, Double Gee or Three-cornered Jack (Spiny Emex), London Rocket, Paradoxa Grass, Stagger Weed and suppression of Wild Radish and Wireweed.	35g	TO AVOID UNACCEPTABLE RESIDUES Do not graze or cut for stock food for 7 weeks after application.

QUEENSLAND ONLY – Post-emergent application

RESTRAINTS

DO NOT apply to crops undersown with legumes.

DO NOT spray when very dry conditions prevail.

DO NOT spray under dry frost conditions.

DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

IF RAIN FALLS within 6 hours of application, the effect could be diminished.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Barley, Wheat – from 3 leaf to early tillering stage	Turnip Weed	6.5g + 300mL Terbutryn®, 500 SC	Spray when weeds are in the 2 to 6 leaf stage, except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.

QUEENSLAND ONLY – Post-emergent application continued

Crop	Weeds Controlled	Rate/ha	Critical Comments
Barley, Wheat – from 3 leaf to early tillering stage	Denseflower Fumitory, Indian Hedge Mustard, Smallflower Fumitory, Wild Turnip	6.5g + 300mL Terbutryn [®] , 500 SC to 10g + 440mL Terbutryn [®] , 500 SC	Spray only when weeds are actively growing. Best results are obtained when good soil moisture has been present since planting.
	Black Bindweed (Climbing Buckwheat), London Rocket	10g + 440mL Terbutryn [®] , 500 SC	Spray only after good rain and when top soil is moist.
Continued	Wild Radish	10g + 440mL Terbutryn [®] , 500 SC to 13g + 600mL Terbutryn [®] , 500 SC	TO AVOID UNACCEPTABLE RESIDUES Do not graze or cut for stock food for 14 days after application.
	Coreopsis, Corn Gromwell (Sheepweed or White Ironweed)	13g + 600mL Terbutryn [®] , 500 SC	

VICTORIA ONLY – Pre-emergent application

RESTRAINTS

DO NOT apply to crops undersown with legumes.

DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Wheat	Annual Ryegrass, Burr Medic, Capeweed, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Volunteer Chickpeas, Faba Beans and Field Peas, Wild Turnip, Wireweed (Hogweed), Yellow Burrweed (Amsinckia), and suppression of Wild Radish and Skeleton Weed.	30 to 35g	Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10cm combine points. Application should not be made to ridged or excessively cloddy soil. Use rates towards the lower end of the range where broadleaf weeds are the major problem. Use the higher rate where Capeweed, Volunteer Chickpeas, Faba Beans and Field Peas, Wild Radish and Wireweed are the problem. 4Farmers Triasulfuron will provide good control of volunteer grain legumes, however a small proportion of plants may survive and require an overspray to eliminate the potential for grain contamination. For best results apply to moist soil when follow up rain is likely to occur within 7-10 days. Use the lower rate (30g/ha) on sandy clay loams with a pH greater than 8.5. For Skeleton Weed a significant degree of control will be achieved on soil types of a predominantly sandy clay loam mixture with a pH greater than 8. Best control is observed where Skeleton Weed germinates in the very early stages of the crop. Surviving plants will be stunted.
	Annual Ryegrass, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Wild Turnip, Wireweed (Hogweed), Ward's weed	10g to 15g plus 1L of a 400g/L trifluralin	This mixture to be only used on alkaline soils where pH is greater than 8 (1:5 Soil:Water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and incorporate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall(greater than 50mm) within 7 days of application may affect efficacy especially at the 10g/ha rate. Late germination of some weeds eg. annual ryegrass will not be controlled in seasons of above average rainfall. Use the higher rate where heavy density of weeds is expected.

VICTORIA ONLY – Post-emergent application

RESTRAINTS

DO NOT apply to crops undersown with legumes.

DO NOT spray when very dry conditions prevail.

DO NOT spray under dry frost conditions.

DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

IF RAIN FALLS within 6 hours of application, the effect could be diminished.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Barley, Oats, Wheat Late Post-emergent application	Wild Radish	15g plus crop oil at 1L per 100L of spray mixture or surfactant at recommended label rates.	Late Post Emergent Application: Spray during early flowering. DO NOT apply to the crop during crop anthesis or flowering (Zadoks 60-69). DO NOT spray on weeds under stress. WARNING: Application to oats may cause some yellowing.
Barley, Oats, Triticale, Wheat – from 3 leaf to early tillering stage	Paterson's Curse, Turnip Weed	6.5g + 300mL Terbutryn [®] , 500 SC	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for field peas, 10 leaflet for vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist.
	Hedge Mustard, Indian Hedge Mustard, Wild Turnip	6.5g + 300mL Terbutryn [®] , 500 SC to 10g + 440mL Terbutryn [®] , 500 SC	Spray only when weeds are actively growing.
	Crassula, Faba Beans (volunteer), Hyssop Loosestrife, Medic, Prickly Lettuce (Whipthistle)	10g + 440mL Terbutryn [®] , 500 SC	Spray only after good rain and when top soil is moist.
	Deadnettle, Field Pea (volunteer), Wild Radish	10g + 440mL Terbutryn [®] , 500 SC to 13g + 600mL Terbutryn [®] , 500 SC	
	Lupins (volunteer), Vetch, Wireweed (less than 3 leaves) suppression.	13g + 600mL Terbutryn [®] , 500 SC	Best results are obtained when good soil moisture has been present since planting.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

PRE-EMERGENT APPLICATION: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION

POST-EMERGENT APPLICATION: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION

HARVEST PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

4Farmers Triasulfuron 750 WG Selective Herbicide is a water dispersible granular herbicide for the pre-plant, incorporated by sowing, control of annual ryegrass, paradoxa grass and certain broadleaf weeds in wheat and for post-emergent control of Wild radish in wheat, oats and barley.

Crops other than wheat, barley, oats, triticale and cereal rye can be very sensitive to low soil concentrations of 4Farmers Triasulfuron 750 WG Selective Herbicide, thus prior to using the product, careful consideration should be given to crop rotation plans.

RESISTANT WEEDS WARNING

GROUP B HERBICIDE

4Farmers Triasulfuron 750 WG Selective Herbicide is a member of the sulfonylurea group of herbicides and has the ALS Inhibitor mode of action. For weed resistance management this is a mode of action Group B herbicide.

Some naturally-occurring weed biotypes resistant to 4Farmers Triasulfuron 750 WG Selective Herbicide and other sulfonylurea herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by 4Farmers Triasulfuron 750 WG Selective Herbicide or other Group B (ALS Inhibitor) herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, 4Farmers Australia Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Triasulfuron 750 WG Selective Herbicide to control the resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a 4Farmers Australia Pty Ltd representative.

MIXING

4Farmers Triasulfuron 750 WG Selective Herbicide is a water dispersible granular herbicide which mixes readily with water

and is applied as a spray.

- Partly fill the spray tank with water
- Start the agitation
- Add the correct amount of product to the spray tank with the agitation system running
- Continue agitation while topping up the tank with water and while spraying
- Use the spray mix within 24 hours of preparation.

APPLICATION

Ground application: Apply by boom spray, applying 30 to 100 litres of water per hectare.

Avoid overlapping of boom runs.

Aerial application: Always spray in a cross wind of less than 5 knots.

Ensure good spray coverage is obtained. Apply 20 to 40 litres per hectare.

SPRAYER CLEANUP

Where the sprayer is being used to spray cereal crops, rinse the sprayer thoroughly with water. Where the sprayer is being used to spray crops other than cereals:

- Drain tank and rinse tank and spray boom with clean water for at least 10 minutes.
- Fill the tank with clean water and add to it 300mL of 4Farmers Spray and Boom Cleaner as per instructions or household chlorine bleach (containing 4% chlorine) per 100L water. Rinse hoses and boom and leave in tank for 15 minutes whilst agitating. Drain through nozzle.
- Repeat step 2 and then rinse thoroughly with clean water to remove all traces of chlorine bleach.
- Nozzles and filters should be cleaned separately.
- Dispose of all water used for cleaning.

COMPATIBILITY

When using a tank mix of 4Farmers Triasulfuron and Spray-Seed*, add 4Farmers Triasulfuron 750 WG, with constant agitation to approximately half the total volume of water to be used.

Ensure that the 4Farmers Triasulfuron is fully dispersed.

Add the Spray-Seed*, fill the spray tank to full volume with water and mix thoroughly.

Apply tank mix immediately, under constant agitation.

4Farmers Triasulfuron 750 WG is compatible with Metolachlor, Tri-alleate, Spray-Seed*, Glyphosate, Sodium molybdate, Zinc sulphate, Manganese sulphate, copper sulphate, 2 4 D Amine, Metsulfuron, Chlorpyrifos, Permethrin, Bromoxynil and Trifluralin.

CROP ROTATION GUIDELINES – Pre-Emergent Application

Where the product is applied at the rate of 30 to 35g/ha:

Unless otherwise specified (see table below), wheat, barley, oats, triticale and cereal rye can be planted the following season without restriction.

For other specified crops the 4Farmers Triasulfuron treated area may be replanted after the interval indicated in the table below. These recommendations are made on the assumption that 4Farmers Triasulfuron is applied to a wheat crop that reaches maturity in the season of application.

Soil pH (1:5 Soil:Water suspension method)	State	Replanting Interval	Minimum rainfall requirements between application and sowing the following crop	Crop
6.5 or less	WA, SA, NSW, ACT, Vic, Qld only	12 months	300mm	Field peas, Linseed, Lucerne, Lupins, Medics**, Subterranean clover**, Faba beans, Chickpeas and Canola.
	NSW, ACT, Qld only	15 months	700mm	Sorghum, Maize, Soybean, Cotton, Cowpea and Mung bean.
6.6 to 7.5	NSW, ACT, Qld only	18 months	900mm	Sunflowers.
		12 months	500mm	Chickpeas and Canola.
		15 months	700mm	Sorghum, Maize, Soybean, Cotton, Cowpea and Mung bean.
6.6 to 7.5	NSW, ACT, Qld only	18 months	900mm	Sunflowers.
		22 months	500mm	Field peas, Linseed, Lucerne, Lupins, Medics**, Subterranean clover**, Faba beans, Chickpeas, Canola, Sorghum, Maize, Soybean and Cotton.
		12 months	250mm	Barley, Oats, Cereal Rye for grain crops.
7.6 to 8.5	Vic, SA only	12 months	300mm	Barley, Oats, Cereal Rye for hay crops.
	NSW, ACT, Qld only	12 months	500mm	Chickpeas and Canola.
	WA, SA, Vic only	18 months	700mm	Sorghum, Maize, Soybean, Cotton, Cowpea and Mung bean.
		24 months	700mm	Field peas, Linseed, Lucerne, Lupins, Medica**, Subterranean clover**, Faba beans, Chickpeas, Canola, Sorghum, Maize, Soybean and Cotton.
8.6 and above	Vic, SA only	12 months	250mm	Barley, Oats, Cereal Rye for grain crops.
		300mm	Barley, Oats, Cereal Rye for hay crops.	
8.6 and above	WA, SA, NSW, ACT, Vic, Qld only	24 months	700mm	Field peas, Linseed, Lucerne, Lupins, Medica**, Subterranean clover**, Faba beans, Chickpeas, Canola, Sorghum, Maize, Soybean and Cotton.

** Includes natural regeneration of subterranean clover and medics. For all other crops seek advice from a 4 Farmers Australia Pty Ltd representative.

Where the product is applied at 10 to 15g/ha plus 1 litre of 400g/L Trifluralin:

Where the pH is less than 7.5 the following crops can be replanted from 9 months after application 4Farmers Triasulfuron providing 300mm of rainfall has been recorded; Field peas, Canola, Chickpeas, Medics, Clover, Lucerne, Safflower, Lupins, Cereal Rye, Barley, Oats, Wheat, Triticale. Where rainfall is less than 300mm for this period further advice should be sought from 4Farmers Australia Pty Ltd.

Where the pH is above 7.5 the following crops can be replanted 9 months after application of 4Farmers Triasulfuron – Cereal Rye, Wheat, Oats, Barley, Triticale.

Where the pH is above 7.5 the following crops can be replanted 12 months after application providing 350mm of rainfall has been recorded; Field Peas, Canola, Chickpeas, Medics, Clover, Lucerne, Safflower, Lupins. Where less than 350mm of rain has fallen between application and sowing the crop, further advice should be sought from 4Farmers Australia Pty Ltd.

CROP ROTATION GUIDELINES

Post Emergent Application

Soil pH (1:5 Soil:Water suspension method)	State	Replanting Interval	Crop
6.5 or less	WA, SA, NSW, ACT, Vic only	7 months	Field peas, Linseed, Lucerne, Lupins, Medics**, Subterranean clover**, Faba beans, Chickpeas and Canola.
		14 months	Sorghum, Maize, Soybean, Cotton.
6.6 to 8.0	WA, SA, NSW, ACT, Vic only	20 months	Chickpeas, Canola, Sorghum, Maize, Soybean, Cotton, Field peas, Linseed, Lucerne, Medics**, Subterranean clover**, Faba beans.

** Includes natural regeneration of subterranean clover and medics. For all other crops a replanting interval of 24 months has to be observed.

These recommendations are made on the assumption that 4Farmers Triasulfuron is applied to a wheat crop that reaches maturity in the season of application.

Where 4Farmers Triasulfuron is used on soil types with pH greater than 8(1:5 Soil:Water suspension test), further advice should be sought from 4Farmers Australia Pty Ltd regarding crop rotation guidelines, except for barley, cereal rye, oats, triticale and wheat.

Crop Rotation Guidelines (tank mixture of 4Farmers Triasulfuron and Terbutryn)

The following crops can be planted after an application of 4Farmers Triasulfuron and Terbutryn:

In areas where pH is less than 7.5

- 9 months** after an application: barley, cereal rye, chickpeas, faba beans, fieldpeas, lupins, medics, oats, canola, safflower, subclover, triticale, wheat.

In areas where pH is greater than 7.5

- 9 months** after an application: barley, cereal rye, oats, triticale, wheat.
- 14 months** after an application: cotton, maize, sorghum, soybeans, sunflowers.

Where residual herbicides are applied following an application, the crop rotational guidelines for these products must be followed.

- 22 months** after an application: chickpeas, faba beans, field peas, lupins,medics, canola, safflower, subclover.

For all other crops advice should be sought from 4Farmers Australia Pty Ltd.

Precaution

Some crop yellowing or crop retardation may occur where stress factors such as water logging, drought, excessive soil acidity or alkalinity, nutrient or trace element deficiency, disease – Rhizoctonia, Take-All, cereal cyst nematodes or soil insects are present or occur following application.

Special care should be taken with regard to the application of 4Farmers Triasulfuron to durum wheats as these may be more sensitive where the above stresses are present. In these situations crop recovery will be rapid provided the stress factors do not continue exerting a negative effect on the crop's growth. Crop retardation may also occur in some instances where considerable late summer/early autumn weed growth occurs. Weeds such as goosefoot, Chenopodium spp can release herbicidally active compounds into the soil.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply on or near shrubs, trees, lawns or crops other than wheat, oats and barley.

DO NOT drain or flush equipment on, or near desirable trees or other plants, where their roots may extend or in situations where by movement of soil, or seepage, absorption of the herbicide may occur.

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift over nearby susceptible plants or crops, cropping lands or pastures.

DO NOT allow spray to drift onto adjacent crops and non-target desirable plants.

DO NOT allow spray to drift onto adjacent fallow land.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original containers in a dry well-ventilated area, as cool as possible out of direct sunlight.

Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. Break, crush,puncture or bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

Re-entry Period

DO NOT enter treated areas without protective clothing until spray has dried.

FIRST AID

f poisoning occurs, contact a doctor or Poisons Information Centre on 13 11 26.

SAFETY DATA SHEET

For further information please refer to the Safety Data Sheet. For a copy visit our website at www.4farmers.com.au

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

4Farmers Australia Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. 4Farmers Australia Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

* Registered Trade Mark

In a Transport Emergency Dial 000 Police or Fire Brigade	Special Poisons Advice: 13 11 26
-----------------------------------------------------------------	-----------------------------------------