

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



TURF CONTROL

Selective Herbicide

ACTIVE CONSTITUENTS:

300 g/L MCPA PRESENT AS THE POTASSIUM SALT
20 g/L CLOPYRALID PRESENT AS THE POTASSIUM SALT
15 g/L DIFLUFENICAN

GROUP F | I HERBICIDE

For the control of Certain Broadleaf Weeds in Turf
as specified in the Direction for Use.

**IMPORTANT: READ THIS LEAFLET
BEFORE USING THIS PRODUCT**

4 FARMERS AUSTRALIA PTY LTD

A.B.N. 51 160 092 428

1 / 70 MCDOWELL STREET, WELSHPOOL, WA 6106

TEL: (08) 9356 3445 www.4farmers.com.au

Batch Number:

Date of Manufacture:

APVMA Approval No: 62327/126277

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT APPLY if weeds or turf are under stress or when frosts are imminent.

DO NOT USE if rain is expected within 4 hours.

DO NOT USE clippings as mulch around other plants or for composting.

DO NOT USE on bent grass or winter grass golf and bowling greens.

Situation	Weeds Controlled	Rate	Critical Comments
Turf	Broadleaf weeds including; White Clover <i>(Trifolium repens)</i> Plantain <i>(Plantago lanceolata)</i> Capeweed <i>(Arctotheca calendula)</i> Cat's ear <i>(Hypochoeris radicata)</i> Bin-dii <i>(Jo-Jo, Onehunga)</i> <i>(Soliva sessilis)</i> Cudweed <i>(Gnaphalium spp.)</i> Creeping oxalis <i>(Oxalis corniculata)</i>	5 L in 200- 500 L water/ha or 50 mL in 2-5 L water per 100m ²	Apply to actively growing weeds. Transient discolouration may occur on kikuyu and carpet grass and Queensland blue couch. Varietal differences in certain buffalo grasses (eg. ST85) may produce more pronounced effects and it is recommended that small areas be tested for turf safety before large-scale application occurs. Avoid exceeding application rates through overlapping sprayed areas. Use a surfactant for difficult to wet weeds.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER,
CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER
APPROPRIATE LEGISLATION.**

WITHHOLDING PERIOD

**DO NOT GRAZE TREATED TURF OR FEED TURF CLIPPINGS FROM
ANY TREATED AREA TO POULTRY OR LIVESTOCK**

GENERAL INSTRUCTIONS

For best results the turf and weeds should be actively growing at the time of treatment. Complete weed death may take up to 6 weeks. Faster results can be expected on smaller weeds. Due to residual effects of diflufenican over-seeding treated areas within 3 months of application may inhibit establishment of the turf grass.

Pourability: If cold temperature environments (less than 5° C) are encountered during product preparation for application the product will be difficult to pour. Should this be the case, it is recommended to shake or stir the contents of the product container for about 5 minutes prior to use.

Mixing: Shake original container before use. Agitate continuously to ensure thorough mixing before and during application. Only mix sufficient chemical for each day's work. Surfactants (if used) should be added after 4Farmers Turf Control Herbicide and water are mixed when the spray-tank is almost full.

Mowing: Avoid mowing during the 3 to 4 days preceding or following treatment.

**In a Transport Emergency
Dial 000
Police or Fire Brigade**

**Special Poisons Advice:
13 11 26**

CLEANING SPRAY EQUIPMENT: THOROUGHLY CLEAN ALL APPLICATION EQUIPMENT AFTER USE. Rinse water should be discharged onto a designated disposal area or if unavailable onto unused land away from desirable plants and water-courses.

Partial Cleaning (before spraying tolerant turf): After using 4Farmers Turf Control Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the tank with a pressure hose. Quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles. Drain and repeat procedure twice.

Complete cleaning (before spraying susceptible plants): After using 4Farmers Turf Control Herbicide empty the tank completely and drain the whole system. Thoroughly wash inside the tank with a pressure hose. Quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles, then drain. Quarter fill the tank again and add an alkaline detergent at 500 ml/100 L water and circulate through the system for at least 15 minutes. Drain, remove filters and nozzles and clean separately. Rinse inside the tank thoroughly using a pressure hose and flush system with clean water.

COMPATIBILITY: No information is available that indicates physical compatibility with other products. Tank-mixing other products with 4Farmers Turf Control Herbicide cannot therefore be recommended. As formulations of other manufacturers' products are beyond the control of 4Farmers Australia Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

RESISTANCE WARNING

GROUP F | I HERBICIDE

4Farmers Turf Control Herbicide is a member of the nicotinilide, phenoxy and pyridine groups of herbicides.

4Farmers Turf Control Herbicide has the process of inhibitors of carotenoid biosynthesis and disruptor of cell growth modes of action. For weed resistance management 4Farmers Turf Control Herbicide is a group F and I herbicide.

Some naturally occurring weed biotypes resistant to 4Farmers Turf Control Herbicide and other Group F and I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by 4Farmers Turf or other Group F or I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, 4Farmers Australia Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Turf Control Herbicide to control resistant weeds.

PRECAUTIONS

Manual Handling of Turf: Hand-weeding and transplanting should not be performed before 2 weeks after spray application, unless workers wear overalls (or equivalent clothing) and chemical resistant gloves.

Re-Entry Statement: Do not allow entry into treated areas until the spray has dried. When prior entry is necessary wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves.

Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Spray drift from 4Farmers Turf Control Herbicide application may cause injury to herbaceous plants, vegetables, fruit crops, vines, ornamentals and trees.

DO NOT use clippings from treated areas for mulch around vegetables or fruit trees.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic life. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

(250ml to 20 L only): Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID INSTRUCTION

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

For further information please refer to the Safety Data Sheet. For a copy visit our website at www.4farmers.com.au

EXCLUSION OF LIABILITY

4Farmers Australia Pty Ltd and/or its affiliates ("4Farmers") shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on 4Farmer's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of 4Farmers Australia Pty Ltd has any authority to add to or alter these conditions.