

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



FLUTRIAFOL 500 SC

Fungicide

ACTIVE CONSTITUENT: 500 g/L FLUTRIAFOL

GROUP 3 FUNGICIDE

For control of certain fungal diseases on wheat, barley and canola when mixed with fertilizer as per Directions for Use.

ALWAYS READ THE LABEL AND USE ONLY AS DIRECTED

4 FARMERS AUSTRALIA PTY LTD

A.B.N. 51 160 092 428
1 / 70 MCDOWELL STREET, WELSHPOOL, WA 6106
TEL: (08) 9356 3445 www.4farmers.com.au

Batch Number:

Date of Manufacture:

APVMA Approval No: 80589/101174

GENERAL INSTRUCTIONS

Fertiliser Application

Granulated Fertiliser: To determine the correct volume of 4Farmers Flutriafol 500 SC required to be applied, refer to Critical Comments in the Directions for Use table, which provides a guide based on a range of common fertiliser application rates. Ensure that the treated fertiliser is dry before sowing with the crop.

Treated fertiliser should be applied in a band 3–4cm below or to the side of seed in planting furrow. It should not be broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered.

Do not mix seed with treated fertiliser and 4Farmers Flutriafol 500 SC.

FUNGICIDE RESISTANCE WARNING

GROUP 3 FUNGICIDE

4Farmers Flutriafol 500 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management 4Farmers Flutriafol 500 SC Fungicide is a Group 3 fungicide. Some naturally occurring individual fungi resistant to 4Farmers Flutriafol 500 SC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by 4Farmers Flutriafol 500 SC Fungicide and other Group 3 fungicides thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use 4Farmers Australia Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Flutriafol 500 SC Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

PRECAUTIONS

Re-entry or Rehandling

Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or

equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility.

If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant local state or territory regulations. Do not burn empty containers or product.

For Refillable Containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

For further information please refer to the Safety Data Sheet. For a copy visit our website at www.4farmers.com.au

CONDITIONS OF SALE

The use of 4Farmers Flutriafol 500 SC Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by 4Farmers Australia Pty Ltd regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4Farmers Australia Pty Ltd accepts no responsibility for any consequences whatsoever resulting from the use of this product.

Not classified as Dangerous Goods by the criteria of the Australian Dangerous Goods Code (ADG Code) for Transport by road or rail.

**In a Transport Emergency
Dial 000
Police or Fire Brigade**

**Special Poisons Advice:
13 11 26**

SEE OVERLEAF FOR DIRECTIONS FOR USE

DIRECTIONS FOR USE: ALL STATES

Crop	Disease	Rate per hectare	WHP	Critical Comments																																							
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	200mL	4 weeks Grazing	<p>Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties.</p> <p>Application: Granulated Fertiliser</p> <p>Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate, which will deposit 200mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide.</p> <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of Granular Fertiliser (mL/100kg)</th> </tr> </thead> <tbody> <tr> <td>60</td> <td>333</td> </tr> <tr> <td>70</td> <td>285</td> </tr> <tr> <td>80</td> <td>250</td> </tr> <tr> <td>100</td> <td>200</td> </tr> <tr> <td>120</td> <td>166</td> </tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of Granular Fertiliser (mL/100kg)	60	333	70	285	80	250	100	200	120	166																											
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of Granular Fertiliser (mL/100kg)																																										
60	333																																										
70	285																																										
80	250																																										
100	200																																										
120	166																																										
Wheat	Stripe Rust (<i>Puccinia striiformis</i> f.sp. <i>tritici</i>)	100– 400mL		<p>For stripe rust control in wheat use 100 to 200mL/ha where up to 100 days protection is required and where heavy stripe rust pressure is expected use 200mL/ha only. Use 300 to 400mL/ha when 120 to 130 days protection is required and where heavy stripe rust pressure is expected, use 400mL/ha only.</p> <p>Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide.</p>																																							
	Septoria tritici blotch (<i>Mycosphaerella graminicola</i>) Take-all (<i>Geaumannomyces graminis</i>)	200mL																																									
Barley	Powdery Mildew (<i>Blumeria graminis</i> f.sp. <i>hordei</i>) Scald (<i>Rhynchosporium secalis</i>)	100mL		<table border="1"> <thead> <tr> <th rowspan="2">Granular Fertiliser rate (kg/ha)</th> <th colspan="4">Volume of product per 100 kg of Granular Fertiliser (mL/100kg)</th> </tr> <tr> <th>100mL rate</th> <th>200mL rate</th> <th>300mL rate</th> <th>400mL rate</th> </tr> </thead> <tbody> <tr> <td>50</td> <td>200†</td> <td>400</td> <td>600</td> <td>800*</td> </tr> <tr> <td>60</td> <td>166†</td> <td>333</td> <td>500</td> <td>667*</td> </tr> <tr> <td>70</td> <td>143†</td> <td>285</td> <td>429</td> <td>571</td> </tr> <tr> <td>80</td> <td>125†</td> <td>250</td> <td>375</td> <td>500</td> </tr> <tr> <td>100</td> <td>100†</td> <td>200†</td> <td>300</td> <td>400</td> </tr> <tr> <td>120</td> <td>88†</td> <td>166†</td> <td>250</td> <td>333</td> </tr> </tbody> </table>	Granular Fertiliser rate (kg/ha)	Volume of product per 100 kg of Granular Fertiliser (mL/100kg)				100mL rate	200mL rate	300mL rate	400mL rate	50	200†	400	600	800*	60	166†	333	500	667*	70	143†	285	429	571	80	125†	250	375	500	100	100†	200†	300	400	120	88†	166†	250	333
			Granular Fertiliser rate (kg/ha)	Volume of product per 100 kg of Granular Fertiliser (mL/100kg)																																							
100mL rate	200mL rate	300mL rate		400mL rate																																							
50	200†	400	600	800*																																							
60	166†	333	500	667*																																							
70	143†	285	429	571																																							
80	125†	250	375	500																																							
100	100†	200†	300	400																																							
120	88†	166†	250	333																																							
				<p>* Application volumes over 600mL per 100kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers.</p> <p>† Where application volume is low, product may be diluted with water to improve coverage.</p>																																							

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

APPLIED IN-FURROW AS A FERTILISER TREATMENT

CANOLA, WHEAT AND BARLEY — DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING.