

In emergency, contact:-

Police and Fire Brigade:

Dial 000

National Poisons Information Centre:

Dial 13 11 26

(from anywhere in Australia)



4 FARMERS AUSTRALIA PTY LTD

A.B.N. 51 160 092 428

1/70 McDOWELL STREET, WELSHPOOL,
WESTERN AUSTRALIA 6106

PH (08) 9356 3445 WWW.4FARMERS.COM.AU

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



CHLORSULFURON
750 WG

Water Dispersible Granules

ACTIVE CONSTITUENT: 750 g/kg CHLORSULFURON

GROUP B HERBICIDE

**For the control of Annual (Wimmera) Ryegrass
and certain broadleaved weeds in winter cereal
crops as per the Directions for Use Table.**

READ THE ATTACHED INFORMATION
THOROUGHLY BEFORE USING THIS PRODUCT.

Batch Number:

APVMA Approval No.: 49836/01

Date of Manufacture:

4 FARMERS AUSTRALIA PTY LTD

A.B.N. 51 160 092 428

1/70 McDOWELL STREET, WELSHPOOL,
WESTERN AUSTRALIA 6106

PH (08) 9356 3445 WWW.4FARMERS.COM.AU

Table of Contents

General Instructions	4
Tables of minimum recommended recropping intervals	6
Spray Preparation	8
Use of Surfactant/Wetting Agent	8
Ground Spraying Equipment	8
Aerial Application	8
Sprayer Cleanup	8
Compatibility	9
Protection of Crops, Native and other Non-Target Plants	9
Protection of Wildlife, Fish, Crustacea and Environment	9
Storage and Disposal	9
Safety Directions	10
First Aid	10
Conditions of Sale	11
Directions for Use	10
Application Method – Pre-Sowing (Incorporated by Sowing)	10-17
Application Method – Post Crop and Weed Emergence	18-23

GENERAL INSTRUCTIONS

1. Before using 4Farmers Chlorsulfuron 750

Careful consideration must be given to the effect of soil pH on chlorsulfuron residues and the effect on subsequent crops. As the soil pH increases the rate of product breakdown decreases. 4Farmers Chlorsulfuron 750 should not be used on soil with a pH of 8.6 (in water) or higher as extended soil residual activity could seriously affect crop rotation options beyond normal intervals.

2. Caution re High pH soil

4Farmers Chlorsulfuron 750 is intended for use on land with a soil pH (in water) of 8.5 or lower.

3. Other Crops

Crops other than wheat, barley, oats, triticale and cereal rye can be extremely sensitive to small concentrations of 4Farmers Chlorsulfuron 750 in soil. See crop rotation recommendations.

4. Resistant Weeds Warning

GROUP	B	HERBICIDE
--------------	----------	------------------

4Farmers Chlorsulfuron 750 is a member of the sulfonylurea group of herbicides.

4Farmers Chlorsulfuron 750 has the acetolacetate synthase inhibitor mode of action. For weed resistance management 4Farmers Chlorsulfuron 750 is a Group B herbicide.

Some naturally occurring weed biotypes resistant to 4Farmers Chlorsulfuron 750 and other acetolacetate synthase inhibitor herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by 4Farmers Chlorsulfuron or other acetolacetate synthase inhibitor herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, 4Farmers Australia Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Chlorsulfuron 750 to control resistant weeds. To prevent, or at least minimise, the risk of resistant weeds occurring, use 4Farmers Chlorsulfuron 750 in tank mixes (if appropriate) and/or rotations with herbicides having different mode of action on the same weed species. Do not allow weed escapes to set seed. Other management/prevention tactics include following, grazing of pasture, pasture manipulation and pastures that are spray topped.

If the presence of sulfonylurea resistant weeds is suspected, 4Farmers Chlorsulfuron 750 or other sulfonylurea herbicides should not be used.

Other strategies to minimise the risk of herbicide resistance are available from your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries Department.

5. Grazing Advice

Avoid grazing 4Farmers Chlorsulfuron 750 treated areas within 24 hours of application to optimise weed control.

A nil withholding period is applicable for grazing 4Farmers Chlorsulfuron 750 treated areas.

6. Crop Safety

DO NOT use in winter cereal crops undersown with legume pasture crops, such as medics and clovers.

DO NOT use on furrow or flood irrigated cereals.

Barley/Oats

DO NOT apply 4Farmers Chlorsulfuron 750 before the two-leaf stage of the crop.

Use of 4Farmers Chlorsulfuron 750 under waterlogged conditions may result in reduced barley yields. However, failure to spray under such conditions may result in an even greater yield reduction through weed competition and weed seed carryover.

DO NOT apply to Stirling Barley.

Wheat

DO NOT apply pre-sowing, incorporated by sowing to varieties Avocet and Durati.

DO NOT apply to varieties Kulin, Miling or Cranbrook.

DO NOT apply pre-sowing incorporated by sowing to variety Banks on soils of pH 5.5 or less.

The variety Vulcan may suffer some yield suppression on acid soils particularly under stressed conditions (e.g. aluminium/manganese toxicity, waterlogging, frosts).

General considerations

DO NOT apply 4Farmers Chlorsulfuron 750 to crops that are stressed by any cause (such as severe weather conditions, drought, waterlogging, excessive soil acidity or soil alkalinity, poor nutrient status, disease, nematode or insect damage) as crop injury may result. When treatment is followed by severe stress (such as drought, prolonged cold, frosty or waterlogged conditions) crop yellowing and growth retardation may occur. Crops normally recover without loss of yield. Disease, nematode or insect damage following application may also result in crop injury.

7. S.A. only

DO NOT apply 4Farmers Chlorsulfuron 750 to light sandy ridges low in organic matter as crop injury may occur.

DO NOT apply other sulfonylurea herbicides following the use of 4Farmers Chlorsulfuron 750.

The use of 4Farmers Chlorsulfuron 750 on early sowing date wheat varieties sown out of season (late) on acid soils may result in some yield suppression under stressed conditions (e.g. waterlogging, frosts, aluminium/manganese toxicity, etc).

8. Crop Rotation

Land previously treated with 4Farmers Chlorsulfuron 750 should not be rotated to crops other than those listed in the table below.

Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas.

4Farmers Chlorsulfuron 750 treated areas may be replanted to any of the specified crops after the interval indicated in the following tables.

Tables of minimum recommended recropping intervals for various combinations of soil pH and crops in different states

N.B. THE TABLE BELOW APPLIES TO ALL STATES

MINIMUM RE-CROPPING INTERVAL (MONTHS AFTER APPLICATION)						
	0	3	6	9	12	15
SOIL pH* 6.5 or less	Triticale, Wheat	Cereal Rye	Oats	Barley	Subterranean Clover** Faba beans, Field pea, Linseed, Lucerne, Lupins, Medics**, Canola (Rapeseed), Safflower	Maize, Sorghum Soybeans, Sunflower

N.B. THE TABLE BELOW APPLIES TO QLD, SA, WA & TAS ONLY.

MINIMUM RE-CROPPING INTERVAL (MONTHS AFTER APPLICATION)						
	0	3	9	15	18	22
RAINFALL REQUIREMENT	-	-	-	Minimum of 700 mm		
Soil pH 6.6 - 7.5	Triticale Wheat	Cereal Rye	Barley, Oats	Japanese Millet, Maize, Panicum Millet, Sorghum, Sunflowers and White French Millet	Cotton, Soybeans	Faba beans, Field Peas, Linseed, Lucerne, Lupins, Medics**, Canola (Rapeseed), Safflower Subterranean Clover**

* Soil pH is to be determined by laboratory analysis using the 1:5, soil: water suspension method.

** Includes natural regeneration of subterranean clover and medics. Land previously treated with 4Farmers should not be rotated to crops other than those listed in the above table. Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas.

N.B. THE TABLE BELOW APPLIES TO QLD, SA, WA & TAS ONLY.

MINIMUM RE-CROPPING INTERVAL (MONTHS AFTER APPLICATION)				
	0	15	18	24 MONTHS OR LONGER
RAINFALL REQUIREMENT	-	Minimum of 700 mm		
Soil pH 7.6 - 8.5	Triticale, Wheat	Japanese Millet, Maize, Panicum Millet, Sorghum, Sunflowers, White French Millet	Barley, Oats, Cereal Rye	ROTATE TO CROPS OTHER THAN CEREALS (SUCH AS LISTED ABOVE) ONLY IF A FIELD TEST STRIP OF ROTATIONAL CROP HAS BEEN SUCCESSFULLY GROWN THROUGH TO MATURITY IN THE PREVIOUS SEASON.
8.6 & above	4Farmers Chlorsulfuron 750 IS NOT RECOMMENDED FOR USE ON SOILS OF pH 8.6 & ABOVE.			

N.B. THE TABLES BELOW APPLY TO NSW & VIC ONLY

MINIMUM RE-CROPPING INTERVAL (MONTHS AFTER APPLICATION)					
	0	3	9	22	26
SOIL pH* 6.6 - 7.5	Triticale, Wheat	Cereal Rye	Barley, Oats	Subterranean Clover** Faba beans, Field Peas, Linseed, Lucerne, Lupins, Medics**, Canola (Rapeseed), Safflower	Maize, Sorghum Soybeans, Sunflower

MINIMUM RE-CROPPING INTERVAL (MONTHS AFTER APPLICATION)			
	0	18	24 MONTHS OR LONGER
SOIL pH* 7.6 - 8.5	Triticale, Wheat	Barley, Cereal Rye, Oats	ROTATE TO CROPS OTHER THAN CEREALS (SUCH AS LISTED ABOVE) ONLY IF A FIELD TEST STRIP OF ROTATIONAL CROP HAS BEEN SUCCESSFULLY GROWN THROUGH TO MATURITY IN THE PREVIOUS SEASON.
8.6 & above	4Farmers Chlorsulfuron 750 IS NOT RECOMMENDED FOR USE ON SOILS OF pH 8.6 & ABOVE.		

9. Spray Preparation

4Farmers Chlorsulfuron 750 is a wettable powder formulation to be blended with water and applied as a spray. Partly fill the boomspray-tank with water. Using a graduated measuring jug, measure the amount of 4Farmers Chlorsulfuron 750 required for the area to be sprayed. Add the correct amount of 4Farmers Chlorsulfuron 750 to the boomspray-tank with the agitation system operating. Fill up to the correct volume with water. The mixture must be kept in suspension at all times by continuous agitation. In tank mixes 4Farmers Chlorsulfuron 750 must be in suspension before adding other herbicides or surfactant.

For application of 4Farmers Chlorsulfuron 750 with liquid fertiliser, slurry the 4Farmers Chlorsulfuron 750 in water, then thoroughly mix the slurry into the liquid fertiliser. Check compatibility before mixing 4Farmers Chlorsulfuron 750 in fertiliser solutions. DO NOT add a surfactant. Where mixed spray solutions have been allowed to stand, thoroughly re-agitate before using.

10. Use of Surfactant/Wetting Agent

For post-emergence application, always use a wetting agent at 100mL/100L of final spray volume. The use of spraying oils is not recommended. NOTE: DO NOT add a surfactant/wetting agent if 4Farmers Chlorsulfuron 750 is tank mixed with another product that already includes a surfactant/wetting agent in the formulation.

11. Ground Spraying Equipment

Use a boom spray properly calibrated to a constant speed and rate of delivery to ensure thorough coverage and a uniform spray pattern. Avoid overlapping and shut off spray booms when starting, turning, slowing or stopping to prevent injury to the crop. Apply a minimum of 30L prepared spray/ha.

12. Aerial Application

Apply a minimum of 20L/ha water. Avoid spraying in still conditions and in winds likely to cause drift onto adjacent sensitive crops. Avoid spraying where drift can affect areas likely to be sown with sensitive crops - see Crop Rotation Recommendations. Turn off spray boom when passing over creeks and dams.

13. Sprayer Cleanup

To avoid subsequent injury to crops other than wheat, triticale, barley, oats or cereal rye, immediately after spraying with 4Farmers Chlorsulfuron 750 thoroughly remove all traces of the product from mixing and spray equipment as follows:

1. Drain tank and then flush tank, boom and hoses with clean water for a minimum of 10 minutes.
2. Fill the tank with clean water and add 300mL of household chlorine bleach (containing 4% chlorine) per 100L of water. Alternatively, use a proprietary spray cleaning agent in accordance with the manufacturers recommendations. Flush through boom and hoses and allow to stand for 15 minutes with agitation engaged. Finally, drain tank.

3. Repeat Step 2.

4. Nozzles and screens should be removed and cleaned separately. To remove traces of chlorine or spray cleaning agent, rinse the tank thoroughly with clean water and flush through hoses and boom.

CAUTION: DO NOT use chlorine or spray cleaning agent with ammonia. All traces of liquid fertiliser containing ammonia, ammonium nitrate or ammonium sulphate must be rinsed with water from the mixing and application equipment before adding chlorine or spray cleaning agent. Failure to do so will release a gas with a musty chlorine odour, which can cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

14. Compatibility

4Farmers Chlorsulfuron 750 is compatible with direct drill herbicides glyphosate and Spray.Seed¹.

4Farmers Chlorsulfuron 750 does not control Wild Oats. However, 4Farmers Chlorsulfuron 750 is compatible with Wild Oat herbicides 4Farmers Tri-allate 400 E.C, Avadex BW3 and Mataven².

4Farmers Chlorsulfuron 750 is compatible with the commonly used broadleaf herbicides Bromoxynil; MCPA 500; Bromoxynil – MCPA; two, 4-D 500; 2, 4-D Ester 800. 4Farmers Chlorsulfuron 750 is compatible with trifluralin 4Farmers Chlorsulfuron 750 is compatible with a range of insecticides including omethoate, dimethoate, cypermethrin, deltamethrin, Sulficid³ and chlorpyrifos.

NOTE: DO NOT add a surfactant/wetting agent when 4Farmers Chlorsulfuron 750 is tank mixed with another product that already has a surfactant/wetting agent in the formulation.

15. Protection of Crops, Native and other Non-Target Plants

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- DO NOT apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots.
- DO NOT apply or allow spray to drift onto adjacent crops and non-target desirable plants, or onto agricultural land scheduled to be planted to other crops as injury to the crop may occur.

16. Protection of Wildlife, Fish, Crustacea and Environment

- DO NOT contaminate any body of water by spraying, cleaning equipment or disposal of waste.

17. Storage and Disposal

- Store in the closed, original container in a dry, well-ventilated area, as cool as possible but out of direct sunlight.
- Keep from contact with fertilisers, insecticides, fungicides and seeds.
- DO NOT re-use container. Crush and bury the empty container in an approved landfill, or bury under at least 500mm of soil in a non-crop, non-pasture area, away from water sources or homes.

18. Safety Directions

Avoid contact with eyes and skin. Do not inhale spray mist. Wash hands with soap and water after use.

19. First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre on 131126. If swallowed and if more than 15 minutes from a hospital induce vomiting, preferably using Ipecac Syrup APF. For further information, refer to the Material Safety Data Sheet.

20. Conditions of Sale

The use of 4Farmers Chlorsulfuron 750 being beyond the control of the manufacturer, no warranty expressed or implied is given by 4 Farmers Australia Pty Ltd regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4 Farmers Australia Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

Registered Trademark of:

- ¹ Syngenta Crop Protection PTY Limited
- ² BASF Australia LTD
- ³ Monsanto Co., USA

DIRECTIONS FOR USE

RESTRAINTS

DO NOT store suspensions of 4Farmers Chlorsulfuron 750 for more than 2 days otherwise significant breakdown will occur?

DO NOT store tank mixes of 4Farmers Chlorsulfuron 750 with other materials for more than 24 hours as product breakdown may occur.

DO NOT spray for post-emergent weed control if rainfall is expected within four (4) hours.

APPLICATION METHOD - PRE-SOWING (INCORPORATED BY SOWING)

CROP	WEEDS CONTROLLED	STATE (S)	RATE g/ha			CRITICAL COMMENTS
			SOIL TYPE			
			Light to Medium Soils		Heavy Soils	
			SOIL pH			
			Less than 7	7-8.5	8.5 or less	
Wheat & Triticale Only	Annual (Wimmera) Ryegrass	NSW, Vic, SA, WA only	20	15 or 20*	20	*Use the higher rate when paddock history suggests a high weed population can be expected. NOTE: Refer to Critical Comments labelled 'All Weeds' below for optimum application timing and conditions.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD – NIL

APPLICATION METHOD - PRE-SOWING (INCORPORATED BY SOWING)

CROP	WEEDS CONTROLLED	STATE(S)	RATE g/ha	CRITICAL COMMENTS
Wheat & Triticale only	Fumitory, Shepherds Purse	NSW, Vic, SA, WA, Tas only	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
	Wireweed/Hogweed, Deadnettle	All States		
	Rough Poppy	NSW, SA, WA, Tas only		
	Stemless Thistle	SA only		
SEE CROP SAFETY PRECAUTIONS	Soursob	NSW, Vic, SA only	15	Apply only to soils with pH 7.5 or above. Apply after majority of soursobs have emerged and leave the soil undisturbed for 1 - 4 weeks prior to cultivation or sowing. The most effective and reliable control is achieved with early post -emergence applications (EPA) after crop and weed emergence.
	Capeweed	NSW, Vic, SA, WA, Tas only	20	On acid soils of pH 5.5 or less this product will give a shorter period of control in wet years.
	Paradoxa Grass Black Bindweed (Climbing Buckwheat)	Qld only	20	Apply to dry soil before the sowing rain. Mechanical incorporation before the sowing rains is not necessary.
	Annual Phalaris	NSW only	20 + 840 mL Trifluralin 480 g ai/L	If possible, spray and incorporate into the soil in one operation. If this is not possible, incorporation should take place within (4) hours of spraying. Delay may cause inferior weed control. Adjust the rate of trifluralin product applied according to the concentration of that product.
	Barley Grass	NSW, Tas only		

RESTRAINTS

DO NOT store suspensions of 4Farmers Chlorsulfuron 750 for more than 2 days otherwise significant breakdown will occur.

DO NOT store tank mixes of 4Farmers Chlorsulfuron 750 with other materials for more than 24 hours as product breakdown may occur.

DO NOT spray for post-emergent weed control if rainfall is expected within four (4) hours.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD - NIL

APPLICATION METHOD - PRE-SOWING (INCORPORATED BY SOWING)

CROP	WEEDS CONTROLLED	STATE(S)	RATE g/ha	CRITICAL COMMENTS
Wheat & Triticale only (Continued)	Indian Hedge Mustard	All States	15	<p>All Weeds: Best weed control is obtained when rainfall or sprinkler irrigation wets the soil to a depth of 5 to 7.5 cm within 4 weeks after application.</p> <p>WA only - Avoid applying to dry, sandy soils as rapid leaching may occur with early seasons rain. SA only - Before using rates greater than 15g/ha on light to medium soils pH of 7.0 to 8.5, seek further advice.</p> <p>Conventionally Sown Crops: On soils of pH, less than 7.0 apply a spray just before sowing or in conjunction with sowing operation. On soils of pH 7.0 or greater, it is not as critical to time the spray just before sowing. Spray onto non-ridged surface free of large clods. Use low profile 10 cm combine points for sowing.</p> <ul style="list-style-type: none"> Sow at speeds of 10 kph or greater. Use light covering harrows at sowing. If applied to dry soil and sowing is to be delayed, incorporate immediately after spraying to prevent losses from wind erosion.
	Wild Turnip, Mouse-ear Chickweed Paterson's Curse (Salvation Jane)	NSW, Vic, SA, WA, Tas only		
	Yellow Burweed/Amsinckia	NSW, Vic, SA, WA only		
	Charlock	Vic, SA, Tas only		
	King Island Melilot	Vic, SA only		
	Storksbill	Vic, SA, WA, Tas only		
	Pimpermels	NSW, Vic, SA, Tas only		
	Lincoln Weed, Common Iceplant, Ball Mustard	SA only		
	Turnip Weed	Qld, SA only		
	Guildford Grass (Onion Grass)	WA only		
Loosestrife	Vic only			
SEE CROP SAFETY PRECAUTIONS	Corn Gromwell (Sheepweed, White Ironweed)	Qld, NSW, Vic, SA, WA only	20	

RESTRAINTS

DO NOT store suspensions of 4Farmers Chlorsulfuron 750 for more than 2 days otherwise significant breakdown will occur.

DO NOT store tank mixes of 4Farmers Chlorsulfuron 750 with other materials for more than 24 hours as product breakdown may occur.

DO NOT spray for post-emergent weed control if rainfall is expected within four (4) hours.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD - NIL

APPLICATION METHOD - PRE-SOWING (INCORPORATED BY SOWING)

CROP	WEEDS CONTROLLED	STATE(S)	RATE g/ha	CRITICAL COMMENTS
Wheat & Triticale only (Continued) SEE CROP SAFETY PRECAUTIONS	Three Cornered Jack, Doublegee (Spiny Emex)	NSW, Vic, SA, WA, only	20	<p>Direct Drilled Crops: Apply tank mixed with either GLYPHOSATE or SPRAY.SEED® in accordance with manufacturers label recommendations.</p> <p>Crops: DO NOT use more than once per season in any one crop.</p>
	Docks	NSW, Vic, SA, WA, Tas only		
	Fat-Hen	NSW, Tas only		
	Cape Tulip	WA only		
	Spear Thistle, Slender Thistle (Shore Thistle)	Tas only		
	Prickly Lettuce (Whip Thistle), Tree Hogweed	Vic, SA only		
	Saffron Thistle (Suppression only)	NSW, Vic, SA, Qld, Tas only		
	Bromegrass (Suppression only)	NSW, Vic, SA, WA, Tas only		
	New Zealand Spinach	Qld only		
	Mintweed, Saltbush, African Turnip Weed, Slender Celery	NSW, Qld only		
				Gives suppression only if population is 20 plants/m or less

RESTRAINTS

DO NOT store suspensions of 4Farmers Chlorsulfuron 750 for more than 2 days otherwise significant breakdown will occur.

DO NOT store tank mixes of 4Farmers Chlorsulfuron 750 with other materials for more than 24 hours as product breakdown may occur.

DO NOT spray for post-emergent weed control if rainfall is expected within four (4) hours.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD - NIL

APPLICATION METHOD – POST CROP AND WEED EMERGENCE

CROP	WEEDS CONTROLLED	STATE (S)	RATE g/ha			CRITICAL COMMENTS
			SOIL TYPE			
			Light to Medium Soils	Heavy Soils		
			SOIL pH			
			Less than 7	7-8.5	8.5 or less	
Wheat Barley, Oats, Triticale, Cereal Rye	Annual (Wimmera) Ryegrass	NSW, Vic, SA, WA only	20 or 25*	15 or 20*	20 or 25*	Use the higher rate under heavy weed pressure. Apply no later than the 3-leaf stage of Annual Ryegrass. Application of 4Farmers Chlorsulfuron 750 to Annual Ryegrass 2 leaf or greater with water volumes less than 50L/ha may result in reduced efficacy.

APPLICATION METHOD – POST CROP AND WEED EMERGENCE

CROP	WEEDS CONTROLLED	STATE(S)	RATE g/ha	CRITICAL COMMENTS
SEE CROP SAFETY PRECAUTIONS	Corn Gromwell (Sheepweed, White Ironweed)	NSW, Vic, SA, WA only	20	Apply at cotyledon to 2 leaf stage. If applied at a later stage only suppression will occur.
	Black Bindweed (Climbing Buckwheat)	Qld, NSW only		Apply from the cotyledon to 2 leaf stage of weed.
	Hoary Cress	Vic, SA, Tas only		Apply when weeds have fully emerged.
	Wild Radish	All States	15 to 20	Annual Ryegrass, Wild Radish & Deadnettle Use the higher rate under heavy weed pressure. Wild Radish - A follow-up spray with a suitable herbicide may be necessary to control subsequent germinations.
	Deadnettle	Qld, NSW, Vic, SA only		
	Mustards	All States	15	All Weeds - Best weed control is obtained when rainfall or sprinkler irrigation wets the soil to a depth of 5 to 7.5 cm within 4 weeks of application. Where treatment is delayed or where weeds are not actively growing due to adverse conditions results may be slow to appear and weeds may be only stunted or suppressed.
	Wild Turnip, Paterson's Curse (Salvation Jane)	NSW, Vic, SA, WA, Tas only		
Yellow Burrweed (Amsinckia)	NSW, Vic, SA, WA only			

RESTRAINTS

DO NOT store suspensions of 4Farmers Chlorsulfuron 750 for more than 2 days otherwise significant breakdown will occur.
DO NOT store tank mixes of 4Farmers Chlorsulfuron 750 with other materials for more than 24 hours as product breakdown may occur.
DO NOT spray for post-emergent weed control if rainfall is expected within four (4) hours.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.
WITHHOLDING PERIOD – NIL

APPLICATION METHOD – POST CROP AND WEED EMERGENCE

CROP	WEEDS CONTROLLED	STATE(S)	RATE g/ha	CRITICAL COMMENTS
Wheat, Barley, Oats, Triticale & Cereal Rye (Continued) SEE CROP SAFETY PRECAUTIONS	Docks Storksbill (Wild Geranium)	Vic, SA, WA, Tas only	15	All Cereals – DO NOT use more than once per season in any crop. Wheat, Triticale and Cereal Rye – Apply after crop emergence when weeds are small and actively growing (ANNUAL RYEGRASS NO MORE THAN 3 leaves. BROADLEAF WEEDS no more than 5cm in height or diameter – for Black Bindweed refer to specific recommendations.)
	Turnip Weed	Qld, NSW, SA only		
	Mouse-ear Chickweed	NSW, Vic, SA, Tas only		
	Charlock, Pimpernels	NSW, Vic, SA, Tas only		
	Ball Mustard	SA only		
	Guildford grass (Onion grass)	WA only		
	Wireweed/Hogweed	All States	20	Apply after majority of Soursobs have emerged. Apply at cotyledon to 2 leaf stage. Apply at the cotyledon to 4 leaf stage
	Soursob (Oxalis pes-caprae)	NSW, Vic, SA, WA only		
	Denseflower Fumitory (Fumaria densiflora)	NSW, Vic, SA, WA, Tas only		
	Rough Poppy	NSW, SA, WA, Tas only		
Mintweed, Saltbush, African Turnip Weed, Slender Celery	NSW, Qld only			

RESTRAINTS

DO NOT store suspensions of 4Farmers Chlorsulfuron 750 for more than 2 days otherwise significant breakdown will occur.

DO NOT store tank mixes of 4Farmers Chlorsulfuron 750 with other materials for more than 24 hours as product breakdown may occur.

DO NOT spray for post-emergent weed control if rainfall is expected within four (4) hours.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD – NIL

APPLICATION METHOD – POST CROP AND WEED EMERGENCE

CROP	WEEDS CONTROLLED	STATE(S)	RATE g/ha	CRITICAL COMMENTS
Wheat, Barley, Oats, Triticale & Cereal Rye (Continued) SEE CROP SAFETY PRECAUTIONS	Staggerweed	Qld NSW, WA, Tas only	20	Barley & Oats - Apply between 2 leaf stage of crop (3 leaf stage -SA only) and early tillering when weeds are small and actively growing (ANNUAL RYEGRASS NO MORE THAN 3 LEAVES: BROADLEAVED WEEDS no more than 5cm in height or diameter – for Black Bindweed refer to specific recommendations). ALWAYS ADD A SURFACTANT
	Shepherd's Purse	NSW, Vic, SA, WA, Tas only		
	Prickly Lettuce (Whip Thistle)	Vic, Tas only		
	Fat Hen	NSW, Tas only		
	Lincoln Weed	SA only		
	Tree Hogweed	Vic only		
	Cape Tulip	WA only		
	New Zealand Spinach	Qld only		
	Spear Thistle	Tas only		
	Matricaria	WA, Tas only		
	Bifora (Carrot Weed)	SA only	25	
	Stemless Thistle	Vic only		

RESTRAINTS

DO NOT store suspensions of 4Farmers Chlorsulfuron 750 for more than 2 days otherwise significant breakdown will occur.

DO NOT store tank mixes of 4Farmers Chlorsulfuron 750 with other materials for more than 24 hours as product breakdown may occur.

DO NOT spray for post-emergent weed control if rainfall is expected within four (4) hours.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD – NIL