

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



TRIADIMEFON 500

FUNGICIDE DRY

ACTIVE CONSTITUENT: 500 g/kg TRIADIMEFON

GROUP C FUNGICIDE

For use as a fertiliser treatment to control leaf and stripe rust;
and suppression of Take All in Wheat and to
control powdery mildew in Barley.

IMPORTANT: READ THIS LEAFLET BEFORE USE

4 FARMERS AUSTRALIA PTY LTD

A.B.N. 51 160 092 428

1 / 70 McDowell Street, Welshpool, WA 6106

Tel: (08) 9356 3445 Fax: (08) 9356 3447

APVMA APPROVAL No: 61547/0107

GENERAL INSTRUCTIONS:

FUNGICIDE RESISTANCE WARNING

Triadimefon is a member of the D.M.I group of **GROUP C FUNGICIDE** fungicides. For fungicide resistance management, 4Farmers Triadimefon 500 Fungicide Dry is a Group C fungicide.

Some naturally occurring individual fungi resistant to 4Farmers Triadimefon 500 Fungicide Dry and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by 4Farmers Triadimefon 500 Fungicide Dry and other Group C fungicides, thus resulting in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, 4 Farmers Australia Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Triadimefon 500 Fungicide Dry to control resistant fungi.

MIXING

4Farmers Triadimefon 500 Fungicide Dry is formulated to be used only by application to granulated fertiliser, and primarily for on-farm application.

Application: Apply product evenly onto the fertiliser either as it is delivered into storage on-farm, or as it is loaded into the fertiliser truck or seeder. Ideally the powder should be metered via an applicator into an auger or elevator, but can also be added to fertiliser in a front end loader bucket as long as the fertiliser is later augered to ensure complete mixing.

To calculate the rate required per tonne of Fertiliser:

Fungicide rate (g/ha) divided by rate of fertiliser (kg/ha) gives the rate in kg fungicide per tonne of fertiliser.

Eg using 80 kg Fertiliser/ha, and wanting 200 gm/ha fungicide - $200/80 = 2.5$ kg/ton.

PROTECTION OF WILDLIFE, FISH, CRUSTACEAN AND ENVIRONMENT

Harmful to fish

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store the unused product in the closed original container in a dry cool, well-ventilated area out of direct sunlight.

Shake bag contents into applicator or onto fertiliser until the bag is empty. Do not dispose of undiluted chemicals on site. Puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers to a depth of 500mm in a disposal specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

DIRECTIONS FOR USE TABLE

APPLIED AS A FERTILISER TREATMENT

Crops	Disease/ Pest	State	Rate	WHP months	Critical Comments
Barley	Powdery mildew	NSW, VIC, SA, WA ONLY	200 g/ha	3	Use where conditions are conducive to very high rates of disease pressure. Also in years when the incidence of disease is expected to be high as forecast by Department of Agriculture or reputable disease forecasters. Regularly monitor crops from about 10-12 weeks after planting to assess the need for follow-up disease control (eg stripe rust) as the residual protection from the fertiliser application declines. See MIXING section for rates and methods of application to fertiliser.
Wheat	Leaf Rust				
	Stripe Rust				
	Suppression of Take-All				

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

- **DO NOT HARVEST FOR 3 MONTHS AFTER APPLICATION**
- **FEED TREATED WITH THIS PRODUCT MUST NOT BE USED FOR ANIMAL CONSUMPTION, POULTRY FEED OR MIXED WITH ANIMAL FEED**
- **DO NOT MIX LEAVES TREATED WITH THIS PRODUCT WITH FEED INTENDED FOR ANIMAL CONSUMPTION**

SAFETY DIRECTIONS

Poisonous if swallowed. May irritate the eyes and skin. Do not inhale dust. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and a disposable dust mask covering mouth and nose.

After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 131126).

SDS

This can be obtained from your supplier or 4 Farmers Australia Pty Ltd at www.4farmers.com.au

CONDITIONS OF SALE

The use of 4 Farmers Triadimefon 500 Fungicide Dry, being beyond the control of the manufacturer, no warranty expressed or implied is given by 4 Farmers Australia Pty Ltd regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4 Farmers Australia Pty Ltd accepts no responsibility for any consequences whatsoever resulting from the use of this product.

**In a Transport Emergency
Dial 000
Police or Fire Brigade**

**Special Poisons Advice:
13 11 26**

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



TRIADIMEFON 500

FUNGICIDE DRY

ACTIVE CONSTITUENT: 500 g/kg TRIADIMEFON

GROUP C FUNGICIDE

For use as a fertiliser treatment to control leaf and stripe rust;
and suppression of Take All in Wheat and to
control powdery mildew in Barley.

IMPORTANT: READ THIS LEAFLET BEFORE USE

4 FARMERS AUSTRALIA PTY LTD

A.B.N. 51 160 092 428

1 / 70 McDowell Street, Welshpool, WA 6106

Tel: (08) 9356 3445 Fax: (08) 9356 3447

APVMA APPROVAL No: 61547/0107

GENERAL INSTRUCTIONS:

FUNGICIDE RESISTANCE WARNING

Triadimefon is a member of the D.M.I group of **GROUP C FUNGICIDE** fungicides. For fungicide resistance management, 4Farmers Triadimefon 500 Fungicide Dry is a Group C fungicide.

Some naturally occurring individual fungi resistant to 4Farmers Triadimefon 500 Fungicide Dry and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by 4Farmers Triadimefon 500 Fungicide Dry and other Group C fungicides, thus resulting in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, 4 Farmers Australia Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Triadimefon 500 Fungicide Dry to control resistant fungi.

MIXING

4Farmers Triadimefon 500 Fungicide Dry is formulated to be used only by application to granulated fertiliser, and primarily for on-farm application. Application: Apply product evenly onto the fertiliser either as it is delivered into storage on-farm, or as it is loaded into the fertiliser truck or seeder. Ideally the powder should be metered via an applicator into an auger or elevator, but can also be added to fertiliser in a front end loader bucket as long as the fertiliser is later augered to ensure complete mixing.

To calculate the rate required per tonne of Fertiliser:

Fungicide rate (g/ha) divided by rate of fertiliser (kg/ha) gives the rate in kg fungicide per tonne of fertiliser.

Eg using 80 kg Fertiliser/ha, and wanting 200 gm/ha fungicide - $200/80 = 2.5$ kg/ton.

DIRECTIONS FOR USE TABLE

APPLIED AS A FERTILISER TREATMENT

Crops	Disease/Pest	State	Rate	WHP months	Critical Comments
Barley	Powdery mildew	NSW, VIC, SA, WA ONLY	200 g/ha	3	Use where conditions are conducive to very high rates of disease pressure. Also in years when the incidence of disease is expected to be high as forecast by Department of Agriculture or reputable disease forecasters. Regularly monitor crops from about 10-12 weeks after planting to assess the need for follow-up disease control (eg stripe rust) as the residual protection from the fertiliser application declines. See MIXING section for rates and methods of application to fertiliser.
Wheat	Leaf Rust				
	Stripe Rust				
	Suppression of Take-All				

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

- **DO NOT HARVEST FOR 3 MONTHS AFTER APPLICATION**
- **FEED TREATED WITH THIS PRODUCT MUST NOT BE USED FOR ANIMAL CONSUMPTION, POULTRY FEED OR MIXED WITH ANIMAL FEED**
- **DO NOT MIX LEAVES TREATED WITH THIS PRODUCT WITH FEED INTENDED FOR ANIMAL CONSUMPTION**