

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



AZOXY CYPRO

Fungicide

ACTIVE CONSTITUENTS: 200 g/L AZOXYSTROBIN
80 g/L CYPROCONAZOLE

GROUP 3 | 11 FUNGICIDE

For the control of certain fungal diseases in Barley, Wheat and Peanuts as specified in the Directions for Use.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

4 FARMERS AUSTRALIA PTY LTD

A.B.N. 51 160 092 428

1 / 70 MCDOWELL STREET, WELSHPOOL, WA 6106

TEL: (08) 9356 3445 www.4farmers.com.au

Batch Number:

Date of Manufacture:

APVMA Approval No: 83297/107919

DIRECTIONS FOR USE

RESTRAINTS

Cereals only:

DO NOT apply more than 2 applications of 4Farmers Azoxy Cypro Fungicide (or any other strobilurin or Group 3 fungicide) in any one season on the same paddock.

DO NOT exceed a combined total of 800mL/ha of 4Farmers Azoxy Cypro Fungicide in any one season on the same paddock.

Crop	Diseases	Rate/ha	Critical Comments
Barley	Leaf Rust (Puccinia hordei) Net Blotch: Net Form (Pyrenophora teres f. teres) Spot Form (Pyrenophora teres f. maculata)	200 mL to 800 mL	Apply from jointing (Z30) and before infection averages 10% on the flag-2 leaf. Ensure thorough coverage of stems and leaves. Use higher rates under high infection pressure or when longer residual protection is required. Lower rates are effective under low disease pressure but have reduced residual effect. Repeat application at early booting stage if necessary. Two applications of the lower rate will generally provide better control than a single application at the higher rate. Addition of ADIGOR® at 2 % v/v improves disease control, especially at the lower rate.
	Powdery Mildew (<i>Blumeria graminis hordei</i>)	400 mL to 800 mL	Spray at the first sign of the disease during the tillering stage. A repeat spray 21 to 28 days later may be required. Ensure thorough coverage of stems and leaves. Higher rates provide longer protection.
Peanuts	Early Leaf Spot (Cercospora arachidicola) Late Leaf Spot (Cercosporidium personatum) Net Blotch (Phoma arachidicola) Rust (Puccinia arachidicola)	500 mL to 750 mL	Apply 1 to 2 applications of 4Farmers Azoxy Cypro Fungicide at 10 to 14 day intervals from first flowering and before disease symptoms appear or when disease pressure is low. Use the low rate when disease pressure is low e.g. early season plantings, and use higher rates when weather conditions favour disease development and/or the crop has a large canopy. When targeting Net Blotch, or when weather conditions are not favourable for application, add a non-ionic surfactant at its recommended use rate. Resistance Management Disease control may be reduced if strains of pathogens less sensitive to 4Farmers Azoxy Cypro Fungicide develop. DO NOT use 4Farmers Azoxy Cypro Fungicide curatively. DO NOT apply more than 1/3 of the total number of fungicide applications per crop as 4Farmers Azoxy Cypro Fungicide. For example, if the total number of fungicides applied to the crop is 9, DO NOT apply more than 3 applications of 4Farmers Azoxy Cypro Fungicide. DO NOT exceed 3 applications of 4Farmers Azoxy Cypro Fungicide per crop. DO NOT apply more than 2 consecutive applications of 4Farmers Azoxy Cypro Fungicide. DO NOT apply more than 5 DMI sprays per season. 4Farmers Azoxy Cypro Fungicide contains a DMI fungicide and should be included in the total number of DMI sprays.
Wheat	Leaf Rust (Puccinia triticina)	400 mL to 800 mL	Apply after flag leaf is 70% emerged and before disease levels reach 1% of flag leaf area. Consider control if disease is greater than 5 to 10% on any lower leaf layer. Use higher rates under high infection pressure or when longer residual protection is required. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.
	Powdery Mildew (Blumeria graminis tritici)		Spray at the first sign of the disease during the tillering stage. A repeat spray 21 to 28 days later may be required. Ensure thorough coverage of stems and leaves. Higher rates provide longer protection.
	Stripe Rust (Puccinia striiformis)		Spray at the first sign of the disease and before 10 to 20% of leaves are infected. A repeat spray 21 to 28 days later may be required. Use higher rates under high infection pressure or when longer residual protection is required.
	Yellow Spot (Pyrenophora tritici-repentis)		Apply once between 70% flag leaf emergence and early flowering. Use higher rates under high infection pressure or where longer residual protection is required. Apply after 70% flag leaf emergence and before disease levels reach 5% on flag leaf. Higher rates give longer residual protection and often better economic returns.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

Barley, Wheat: Grazing: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 21 DAYS AFTER APPLICATION

Harvest: DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION

Peanuts: Grazing: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 14 DAYS AFTER APPLICATION

Harvest: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION

EXPORT SLAUGHTER INTERVAL

DO NOT slaughter animals destined for export within 7 days of consumption of treated cereal forage or straw, or peanut forage or fodder.

GENERAL INSTRUCTIONS

MIXING

4Farmers Azoxy Cypro Fungicide is a suspension concentrate (SC) formulation that mixes readily with water and is applied as a spray.

1. Partly fill the spray tank with water
2. Start the agitation
3. Add the correct amount of product to the spray tank with the agitation system running
4. Continue agitation while topping up the tank with water and while spraying
5. Use the spray mix within 24 hours of preparation

GROWING CONDITIONS

In general, 4Farmers Azoxy Cypro Fungicide should be applied under good growing conditions with adequate soil moisture. 4Farmers Azoxy Cypro Fungicide should not be applied to crops that are under stress due to very dry, very wet, frosty conditions, nutrient deficiency or high insect pressure, as this may give less than reliable results.

SPRAY TIMING

Always inspect crops to assess disease development immediately before spraying. Best results will be achieved from applications made in the earliest stages of disease development or as a protectant treatment following a disease risk assessment.

Cereals: Applications of 4Farmers Azoxy Cypro Fungicide should be made within the crop growth stages ranging from tillering to early flowering (Z26 to Z62). The object of spraying is to keep the upper 2 to 3 leaves green and functioning through grain filling stage.

Peanuts: The object of spraying is to control diseases and stop defoliation during pegging or pod fill stages to ensure stems are sufficiently strong for harvesting.

APPLICATION

May be applied by boom spray or aircraft. Ensure complete coverage of all leaves and stem is obtained.

Cereals:

Boom Application: As a guide, apply in spray volumes ranging from 50L to 100L water per hectare. The use of flat fan nozzles is recommended using a fine to medium quality spray pattern, with pressures in the range 100kPa to 400kPa.

Aerial Application: As a guide, apply using 20L to 30L water per hectare using a fine to medium quality spray. Use the lower rate when applications are made with a cross wind of not less than 5 knots. Use the higher rates when applying to dense crops.

Peanuts:

Boom Application: As a guide, apply in spray

volumes ranging from 150L to 500L water per hectare. The use of hollow cone nozzles or flat fans is recommended using a fine to medium quality spray pattern, with pressures in the range of 500kPa to 1000kPa for the hollow cones and 150kPa to 300kPa for flat fan nozzles.

Aerial Application: As a guide, apply 25L to 60L water per hectare using a fine to medium quality spray. Use with the lower rate when applications are made with a cross wind of not less than 5 knots. Use the higher rates when applying to dense crops.

COMPATIBILITY/TANK MIXING

4Farmers Azoxy Cypro Fungicide is physically compatible with a wide range of products, however the biological compatibility of these mixtures may not have been fully tested under all environmental and biological conditions.

A mixture of 4Farmers Azoxy Cypro Fungicide with any other product may in some circumstances be ineffective or may cause damage.

If tank mixes are to be used observe all directions, precautions and limitations on all products to be used.

As formulations of other manufacturer's products are beyond the control of 4Farmers Australia Pty Ltd, and the quality of water may vary with location, all mixtures should be tested prior to mixing commercial quantities.

FUNGICIDE RESISTANCE WARNING

GROUP 3 | 11 FUNGICIDE

4Farmers Azoxy Cypro Fungicide is a combination of a DMI and a strobilurin and related products fungicide. For fungicide resistance management 4Farmers Azoxy Cypro Fungicide is both a Group 3 and a Group 11 fungicide.

Some naturally occurring individual fungi resistant to 4Farmers Azoxy Cypro Fungicide and other Group 3 and/or Group 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by 4Farmers Azoxy Cypro Fungicide or other Group 3 and/or Group 11 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, 4Farmers Australia Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Azoxy Cypro Fungicide to control resistant fungi.

PRECAUTIONS

Re-entry period: DO NOT allow entry into treated areas until the spray has dried unless wearing cotton overalls (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

HIGHLY TOXIC TO AQUATIC LIFE. DO NOT contaminate dams, waterways or drains with the chemical or used containers.

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift on adjacent areas, particularly wetlands, waterbodies or watercourses.

DO NOT apply 4Farmers Azoxy Cypro Fungicide aerially if waterbodies, watercourses or wetlands are within 50 metres downwind of the application area.

PROTECTION OF CROPS

WARNING

Azoxystrobin is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees.

DO NOT spray 4Farmers Azoxy Cypro Fungicide where spray drift may reach apple trees.

DO NOT spray when conditions favour drift beyond the area intended for application. Conditions which may contribute to drift include thermal inversions, excessive wind speed, certain spray nozzle/pressure combinations, small spray droplet size, etc.

DO NOT use spray equipment which has been previously used to apply 4Farmers Azoxy Cypro Fungicide to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity.

STORAGE AND DISPOSAL

Store in closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed.

Will irritate the eyes. Avoid contact with eyes.

When opening the container and preparing spray wear:

- Cotton overalls buttoned to the neck and wrist (or equivalent clothing)
- A washable hat
- Elbow-length PVC gloves
- Goggles

Wash hands after use.

After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

For further information please refer to the Safety Data Sheet. For a copy visit our website at www.4farmers.com.au

CONDITIONS OF SALE

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. 4Farmers' liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which 4Farmers Australia Pty Ltd shall have no liability.

In a Transport Emergency Dial 000 Police or Fire Brigade	Special Poisons Advice: 13 11 26
---	---