

# Fluroxypyr 200

Similar to: *Starane*<sup>®</sup>

<b>Active Constituent</b>	Fluroxypyr as Methylheptyl ester 200g/L		<b>Formulation</b>	EC
<b>Typical Situations</b>	Wheat, barley, oats, triticale, winter and summer fallows (see registration section).			
<b>Chemical Group</b>	I	<b>Mode of Action</b>	Foliar and root uptake, readily translocated – causes unregulated, disrupted growth.	
<b>Indicative Price ex GST</b>	\$12.60/L	<b>Typical Pack Size</b>	20L	
<b>Poison Schedule</b>	5	<b>Dangerous Goods Class</b>	-	<b>UN</b> -
<b>Withholding Period</b>	<b>Harvest</b> Not required when used as directed (except Poppies – 10 weeks pre-harvest). <b>Grazing</b> Do not graze or cut for hay for 7 days post application.			
<b>Plant Back</b>	Cereals - 7 days, soybeans - 14 days.			
<b>Application Method</b>	Boom spray greater than 50L/ha, aerially 35L/ha.		<b>Rain fastness</b>	1 hour
<b>Efficacy</b>	Do not spray on stressed weeds.			
<b>Adjuvants</b>	Non-ionic surfactant and crop oils (see label).			
<b>Compatibility</b>	2, 4-D Amine, 2, 4-DB, Atrazine, Clodinafop, Glyphosate, Clopyralid, Diclofop-methyl, MCPA, Metsulfuron-methyl, Picloram and Triclopyr.			
<b>Incompatibility</b>	Follow guidelines for Atrazine, Clodinafop, and Glyphosate.			
<b>Water Quality</b>	Hard water and saline water will effect efficacy			
<b>Time to Effects and Symptoms</b>	3-4 hours – twisting and downward bending. 3 weeks – plant death.			
<b>4F Broadacre Registrations</b>	Wheat, barley, oats and triticale, and summer and winter fallows (extract listed).			
<b>4F Other Registrations</b>	Woody weeds in non-crop, industrial and rights of way and other (see label).			
<b>Similar Product Registrations</b>	As above.			
<b>Situation</b>	<b>Target Weed/s</b>	<b>Rate/ha</b>	<b>Comments</b>	
Winter Cereals <b>post emergent</b> (3 leaf to flag leaf Z13-Z39)	Bedstraw (1-3 whorls), Prickly Lettuce (2-5leaf)	1L	Check application timing of mixing products.	
	Doublegee, Wireweed (2-4 leaf), Volunteer lupins (2-8 leaf)	1.5L	Slight delay in crop timing may occur. Atrazine mixes available on label.	
	Mustards, Shepherd's Purse, Turnip Weed, Wild Radish, Wild Turnip, Bittercress (up to 8 leaf)	1.5L + 700ml LVE MCPA or 1L of MCPA 500	Use with crop oil or wetter. Any lupins present will be severely damaged.	
<b>Comments</b>				
Commonly used as a spike for Glyphosate (@ 400ml/ha). Agitation must be constant and vigorous throughout application. Elapsed time between mixing and use should not exceed 2 hours. Crop oil cannot be used in mixes with Atrazine, if adjuvant is required add Wetter 1000. Performs well with lower water rates and cold water.				