

# Tribenuron Methyl 750

Similar to: *Express*<sup>®</sup>

<b>Active Constituent</b> Tribenuron Methyl 750g/kg		<b>Formulation</b> WG	
<b>Typical Situations</b> Fallow and prior to sowing (see plant back and registration section for timing).			
<b>Chemical Group</b> B	<b>Mode of Action</b> Foliar and root uptake, rapid upward translocation – inhibits cell division and disrupts growth.		
<b>Indicative Price ex GST</b> N/A	<b>Typical Pack Size</b>		2kg
<b>Poison Schedule</b> 5	<b>Dangerous Goods Class</b> -	<b>UN</b> -	<b>Packing Code-</b>
<b>Withholding Period</b>	<b>Harvest</b> Not required when used as directed.		
	<b>Grazing</b> Not required when used as directed.		
<b>Plant Back</b> Wheat, barley and oats - 3 days. Maize, Mung beans, Sorghum and Soybeans -21 days.			
<b>Application Method</b>	Boom spray greater than 50L/ha, aerially 20-30L/ha		<b>Rain fastness</b> 2 hours
<b>Efficacy</b> Apply to young, actively growing weeds in warm, moist conditions to maximise efficacy.			
<b>Adjuvants</b> 0.1% non-ionic wetter (Wetter 1000), add 1% mineral spray oil (Turbo Charge).			
<b>Compatibility</b> Atrazine, Bipyridyls and Glyphosate.			
<b>Incompatibility</b> Grass selectives.			
<b>Water Quality</b> Hard water may affect efficacy.			
<b>Time to Effects and Symptoms</b> 1-3 weeks - plant discolouration followed by death.			
<b>4F Broadacre Registrations</b> Fallow and prior to sowing (extract listed).			
<b>4F Other Registrations</b> N/A			
<b>Similar Product Registrations</b> As above.			
<b>Situation</b>	<b>Target Weed/s</b>	<b>Rate/ha</b>	<b>Comments</b>
Fallow or Prior to crop	New Zealand Spinach (up to 10 leaf), Turnip Weed (until flowering)	20g	For more advanced weeds and/or large populations use the mix recommended below.
	Amaranthus, Caltrop, Common Sowthistle (up to 10 leaf), Deadnettle (up to 6 leaf)	25g	
	Medics, Prickly Lettuce (up to 4 leaf or 50mm diameter)	30g	
Fallow or Prior to crop	Mintweed (up to 10 leaf)	15g + 600ml Glyphosate 450	Often used as a spike with Glyphosate to improve efficacy. Always use a Wetter (non-ionic).
	Common Thornapple (8 leaf), Pigweed (10 leaf), Prickly Lettuce (6 leaf)	20g + 600ml Glyphosate 450	
	Black Bindweed, Deadnettle (10 leaf)	25g + 600ml Glyphosate 450	
<b>Comments</b>			
Do not use if a temperature inversion is likely.			
Do not apply on stressed plants.			
For resistance management, do not use Tribenuron Methyl or another Group B herbicide in consecutive seasons.			
Mixing with Glyphosate may assist with resistance management.			
If heavily grazed, remove stock to allow some regrowth prior to application.			