

Bromacil 800

Similar to: *Uragan*[®]

| | | | | |
|--|--|------------------------------|---|----------------------|
| Active Constituent | Bromacil 800g/kg | | Formulation | WG |
| Typical Situations | Horticultural crops, around agricultural buildings, industrial, commercial, rights of way and other (see registration section). | | | |
| Chemical Group | C | Mode of Action | Root and some foliar uptake, upward translocation – disrupts photosynthesis, destroys cells. | |
| Price ex GST | \$40.00/kg | Typical Pack Size | 5kg | |
| Poison Schedule | N/S* | Dangerous Goods Class | - | UN - |
| Withholding Period | Harvest | N/A | | |
| | Grazing | N/A | | |
| Plant Back | 3 years, everything except pineapples. | | | |
| Application Method | Boom spray 50L/ha minimum, best at 250-450L/ha | | | Rain fastness |
| | | | | N/A |
| Efficacy | Sufficient moisture to carry the chemical into the soil is required within 3-4 days for best efficacy. | | | |
| Adjuvants | Wetter 1000 (if weeds present). | | | |
| Compatibility | Most herbicides (confirm individual mixes using a jar test). | | | |
| Incompatibility | Grass selectives, Group B herbicides, Simazine. | | | |
| Water Quality | - | | | |
| Time to Effects and Symptoms | 7-10 days - chlorosis and desiccation in sunny conditions. | | | |
| 4F Broadacre Registrations | N/A | | | |
| 4F Other Registrations | Citrus, asparagus, pineapple, industrial and rights of way and other (extract listed). | | | |
| Similar Product Registrations | <i>Uragan</i> [®] is registered for fence lines. | | | |
| Situation | Target Weed/s | Rate/ha | Comments | |
| Citrus (established groves only) | Most annual weeds | 2.2 - 4.5kg | Vary rate with soil type (e.g. light soil 2.2kg, heavy soil 4.5kg). Apply to a weed free ground in Autumn to Spring. | |
| | Perennial grasses, e.g. Couch grass, Nut grass, Johnson grass, Kikuyu, Skeleton weed | 4.5 - 6.5kg | | |
| Industrial Areas (around sheds, etc.) | Barley grass, Bathurst Burr, Brome grass, Caltrop, Capeweed, Cobblers peg, Doublegee, Paterson's Curse, Prairie grass, Annual Ragweed, Annual Ryegrass, Saffron thistle, Summer grass, Thornapple, Wild Mustard, Wild oats, Wild Radish, Wild Turnip, Winter grass and other (see label) | 3.5 - 6.5kg | Lower rates are for lighter soils, higher rates for heavier soils. Rates will also increase with increase in rainfall with the lowest rate for 250mm or less per year. Apply retreatment when grasses reappear. Use Wetter 1000 where weeds are present. | |
| | Bahia grass, Bluetop, Guinea grass, Horehound, Para grass, Purple Top, Plantain, St John's Wort, Soursob | 8 - 13kg | | |
| | Bindweed, Couch grass, Dock, Johnson grass, Kikuyu, Nut grass, Skeleton weed, Sorrel, Water Couch | 16.5 - 22kg | | |
| | Retreatment for above | 2 - 6.5kg | | |
| Comments | | | | |
| Do not apply more than 9kg/ha per year in citrus and pineapple crops. In low rainfall environments deep rooted perennials may not be controlled. Commonly used in combination with Paraquat for Glyphosate-resistant weed control on firebreaks and other non-agricultural areas, and provides long periods of soil residual activity. *N/S – Not scheduled | | | | |