

Clopyralid 300

Similar to: *Lontrel*[®]

Active Constituent	Clopyralid 300g/L		Formulation	SL
Typical Situations	Wheat, barley, canola, oats, triticale and pasture (see registration section).			
Chemical Group	I	Mode of Action	Foliar and root uptake, readily translocated – causes unregulated, disrupted growth.	
Price ex GST	\$14.35/L	Typical Pack Size	20L, 110L, 1000L	
Poison Schedule	5	Dangerous Goods Class	-	UN -
Packing Code-				
Withholding Period	Harvest	Cereals - apply at least 10 weeks prior to harvest, canola - apply at least 12 weeks prior to harvest.		
	Grazing	Do not graze or cut for hay for 7 days post application.		
Plant Back	Up to 24 months for broadleaf crop and pasture species (field peas and faba beans are particularly susceptible), 1 week for cereals and canola.			
Application Method	Boom spray greater than 50L/ha, aerially greater than 20L/ha.		Rain fastness	3 hours
Efficacy	Do not spray on stressed weeds.			
Adjuvants	None Required			
Compatibility	Bromoxynil, grass selectives, Phenoxys, SU herbicides, Triazines and Ureas.			
Incompatibility	Trace elements (copper sulphate and zinc sulphate).			
Water Quality	Hard and alkaline water reduces efficacy.			
Time to Effects and Symptoms	Dependant on humidity; varies from hours to weeks – shoot and root inhibition, thickening roots, twisted or curled shoots, cupped or crinkled leaves.			
4F Broadacre Registrations	Wheat, barley, canola, oats, triticale, fallow and pasture (extract listed).			
4F Other Registrations	Forests, rights of way and industrial (see label).			
Similar Product Registrations	As above.			
Situation	Target Weed/s	Rate/ha	Comments	
Winter Cereals (pre-sowing)	Capeweed (up to 8 leaf or 100mm diameter)	150ml+ Knock down herbicide	WA only. Should be used with Glyphosate or Brown Out.	
Winter Cereals (post-sowing, from 2 leaf to jointing Z12-Z35)	Capeweed	150ml	Weeds should be actively growing and not larger than 50mm diameter. Post-sowing pre-emergent through to 3 leaf - can add 300ml/ha Diuron for control of transplants.	
Winter Cereals (4 to 5 leaf onwards Z14)	Capeweed, Soldier thistle (up to 100mm diameter)	300ml	Hairy leaved medics may require spray oil. Weeds may only be stunted but uncompetitive.	
	Volunteer Chickpeas, Lentils (to 6 leaf)	250ml		
	Volunteer Faba Beans, Lupins (to 4 leaf)			
	Volunteer Field Peas (up to 6 nodes)	150ml		
	Volunteer Medics, seedling Lucerne (to 8 leaf), volunteer Sub-clover (to 6 leaf)	150ml		
Comments				
Do not apply Clopyralid on crops intended for export hay.				
Do not apply to pastures/crops intended for compost/mulches. Can damage susceptible crops.				
Decay of Clopyralid is highly effected by rainfall following application, and soil texture – requires at least one 25ml rainfall event & 1 week of residual soil moisture to allow microbial breakdown prior to sowing susceptible species.				
Applications 150ml/ha or greater post-June may substantially impact regeneration of Serradella stands in the following year.				
Anecdotal evidence suggests early applications on young Fleabane may provide suppression or control.				