

## Metaldehyde Snail and Slug bait

|   |  |  |   |                    |
|---|--|--|---|--------------------|
| <b>Active Constituent</b>   | Metaldehyde 15g/kg   |  | <b>Formulation</b>  | Spreadable Pellets |
| <b>Price ex GST</b>   | \$1.32/kg  | <b>Typical Pack Size</b>                         | 20kg, 500kg, 1,000kg  |                    |
| <b>Poison Schedule</b>  | 5  | <b>Dangerous Goods Class</b>                     | -   | <b>UN</b> -        |
| <b>Withholding Period</b>   | <b>Harvest</b>   | Do not apply later than 7 days prior to harvest. |   |                    |
|   | <b>Grazing</b>   | Do not allow stock to graze in treated areas.    |   |                    |
| <b>Efficacy</b>   | Scatter product evenly around shelter sites, when adult snails and slugs are most active (preferably immediately after rain, or during humid, overcast weather, and when green plant material is limited). Late summer and early autumn rains provide ideal application windows. |  |   |                    |
| <b>Compatibility</b>  | N/A  |  |   |                    |
| <b>Water Quality</b>  | N/A  |  |   |                    |
| <b>Situation</b>  | <b>Target Pest/s</b>   | <b>Rate</b>                                      | <b>Comments</b>   |                    |
| Crops, borders and shelter areas  | Snails, Slugs  | 50g/m <sup>2</sup><br>(500kg/ha)                 | Scatter the pellets around the area to be protected and/or where snails and slugs congregate. Spread pellets evenly.  |                    |
| Winter cereals, oilseeds  | Common white snail ( <i>Ceruella virgata</i> ), White Italian snail ( <i>Theba pisana</i> ), Conical snail ( <i>Cochlicella acuta</i> ), Grey filed slug ( <i>Deroceras reticulatum</i> )  | 5 – 7.5 kg/ha                                    | Broadcast pellets evenly over the crop or pasture area to be protected from snails and slugs. Pay particular attention to areas adjacent to headlands, fence lines and other areas where snails and slugs shelter and move into crops or pastures. Use the higher rate where there are large populations of snails or slugs and where wet, cool climatic conditions are favourable for snail and slug movement. |                    |
| Pulses*, Pasture establishment*   |  |  | *Pulse and pasture crops only: This product is only to be used during the establishment stages of pulse crops and pastures. Broadcast pellets evenly over the crop area to be protected prior to planting, post-plant pre-emergence or early post-emergence (prior to 4 leaf stage).  |                    |
| <b>Comments</b>   |  |  |   |                    |
| <p><b>CARE:</b> - This product is poisonous, avoid inhalation and contact with skin and eyes.</p> <ul style="list-style-type: none"> <li>- Keep domestic animals and livestock away from treated areas.</li> <li>- <b>Do not</b> apply to edible plants.</li> </ul> <p>Metaldehyde Snail and Slug bait is a spreadable pelletized poison for the broad acre control of snails and slugs. Product breaks down in the soil over time leaving no harmful residues. May require re-application in problem areas.</p> <p>A high number of bait points on the ground is critical for good snail control. 4Farmers pellets contain 15g/kg Metaldehyde and have an ideal smaller size. Spreading less than 20kg/ha will economically achieve around 45 pellets/m<sup>2</sup>.</p> |  |  |   |                    |