

Sulfosulfuron 750

Similar to: *Monza*[®]

Active Constituent	Sulfosulfuron 750g/kg		Formulation	WG
Typical Situations	Wheat and triticale.			
Chemical Group	B	Mode of Action	Root and foliar uptake, rapid upward translocation – inhibits cell division and disrupts growth.	
Price ex GST	\$224.00/kg	Typical Pack Size	500g	
Poison Schedule	N/A	Dangerous Goods Class	-	UN -
Withholding Period	Harvest	Not required when used as directed.		
	Grazing	Not required when used as directed.		
Plant Back	For all non-label crops: 12 months for soil pH below 6.5, 22 months for soil above pH 6.5.			
Application Method	Boom spray 40-100L/ha, do not apply aerially			Rain fastness 1 hour
Efficacy	Incorporate immediately after pre-sowing application. Moist soil (50-75mm deep) is required to activate product (within 10 days for optimum efficacy). Post-emergent control is maximised in conditions where weeds are young and actively growing.			
Adjuvants	Always add paraffin oil (e.g. Take Up) when applying post-emergent.			
Compatibility	Clopyralid, Glyphosate, LVE MCPA, Paraquat, Tri-Allate and Trifluralin.			
Incompatibility	Dicamba, Diclofop-Methyl, Terbutryn and Tralkoxydim, Dimethoate and Omethoate.			
Water Quality	N/A			
Time to Effects and Symptoms	Pre-sowing application: Any emerging weeds will be stunted and uncompetitive. Post emergent application: 1-4 days - Chlorosis then death.			
4F Broadacre Registrations	Wheat and triticale (extract listed).			
4F Other Registrations	-			
Similar Product Registrations	As above.			
Situation	Target Weed/s	Rate/ha	Comments	
Wheat, Triticale (pre-emergent)	Barley Grass, Brome grass, Wild Turnip	25g	Apply to bare soil pre/at sowing for incorporation by sowing.	
	Above + Wild Oats	25g + 1.6L Tri-Allate 500 EC	In no-till systems apply with Glyphosate.	
	All above + Annual Ryegrass, Canary Grass, Fumitory, Wireweed	25g + 800ml Trifluralin 480 EC	For best efficacy when mixing, follow incorporation instructions of additive.	
Wheat, Triticale (post-emergent, first leaf to early tillering Z11-22)	Amsinckia, Wild Mustard	Cotyledon-4 leaf	25g	Optimum results will be obtained by good soil moisture at and post application. Follow-up spray using a different herbicide is required for subsequent germinations.
	Barley Grass, Silver Grass, Wild Oats	Z11 - Z14		
	Brome grass	Z11 - Z13	25g	
	Volunteer Canola, Wild Radish, Wild Turnip	Cotyledon to 4 leaf	20g	
	Volunteer Field Peas	1-3 leaf	20g	
Comments				
Do not use this product sequentially (within the year) to avoid grain contamination.				
Do not apply to crops under sown with a legume.				
Do not spray when very dry conditions prevail.				
Do not use on furrow or flood irrigated crops.				
Do not apply to soils with pH over 8.5.				
Do not use if another SU herbicide was used in the same season.				
Do not store in direct sunlight for long periods of time.				
CARE: Product affects many species even at low rates of application. Beware of residual effects on following crops.				