

**FOR ANIMAL TREATMENT ONLY
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



DICYCLANIL 50 g/L

SPRAY-ON FLY TREATMENT

ACTIVE CONSTITUENT: 50 g/L DICYCLANIL

For the protection of sheep, either off-shears or with any length wool, against fly strike (caused by dicyclanil-susceptible strains of blowflies (*Lucilia cuprina*)) for 18 to 24 weeks.

For the protection of mulesing and marking wounds on sheep against fly strike (caused by dicyclanil-susceptible strains of blowflies (*Lucilia cuprina*)) during the wound healing process.

4Farmers Dicyclanil 50 g/L Spray-On Fly Treatment contains dicyclanil, a member of the pyrimidine group of chemicals. It is effective against strains of *Lucilia cuprina* that are sensitive to dicyclanil. It is not as effective against strains of *Lucilia cuprina* that develop resistance to dicyclanil. Resistance may develop to any chemical. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance.

**IMPORTANT: READ THE ATTACHED INFORMATION
THOROUGHLY BEFORE USING THIS PRODUCT**

NET CONTENTS: 20L

4 FARMERS AUSTRALIA LIMITED
ABN: 51 160 092 428
70 MCDOWELL ST, WELSHPOOL
TEL: 1800 038 445
WWW.4FARMERS.COM.AU

APVMA APPROVAL NO.: 92570/135915



DICYCLANIL 50 g/L SPRAY-ON FLY TREATMENT



DIRECTIONS FOR USE

RESTRAINTS

DO NOT use on ewes which are producing or may in the future produce milk that may be used or processed for human consumption. DO NOT re-treat animals for a minimum of 9 weeks after last treatment. Side Effects: NIL

DOSAGE AND ADMINISTRATION

4Farmers Dicyclanil 50g/L Spray-On is applied as a low-volume treatment onto areas where protection is required. When applying 4Farmers Dicyclanil 50g/L Spray-On the nozzle of the applicator gun should be high enough above the fleece (approximately 15-25 cm), to achieve a 15 cm wide band at each pass for protection of body strike. For treatment of mulesing wounds, the band must be wide enough to cover the wound and 25mm onto the surrounding wool. DO NOT apply during gusty or windy weather as spray drift may reduce the period of protection. It is good agricultural practice for mulesing/ marking to be carried out at times of minimal fly activity. Reduced efficacy may occur in sheep affected with dermatophilosis.

GENERAL DIRECTIONS

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER. CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

4Farmers Dicyclanil 50g/L Spray-On provides season long protection to sheep against body, breech and poll strike and may be used either off-shears (within 24 hours of shearing) or on sheep with any length wool (Short Wool: greater than one day and up to 6 weeks after shearing; Long Wool: greater than 6 weeks after shearing).

4Farmers Dicyclanil 50g/L Spray-On also protects mulesing and marking wounds on lambs against fly strike during the wound healing process. The dose volume is determined by the weight of sheep being treated. Apply Neovek Spray-On according to the diagrams and tables in the Directions for Use. DO NOT dilute. DO NOT mix with any other product. Apply directly from the container or an appropriate back pack using a spray-on applicator gun. 4Farmers Dicyclanil 50g/L Spray-On may be applied to all breeds of sheep to obtain protection against fly strike.

Significant soiling with dags, mud, dust or urine may reduce efficacy. Where significant dags are present, a light crutching is indicated prior to treatment. Protection period may be reduced if sheep scour heavily following application.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 28 days before slaughter for human consumption.

MILK: DO NOT USE on sheep which are producing or may in the future produce milk that may be used or processed for human consumption.

WOOL HARVESTING INTERVAL: DO NOT USE less than 3 months before shearing or fibre collection.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 120 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the registrant (02) 9264 3210 or APVMA website (www.apvma.gov.au/residues).

Use of this product may result in wool residues that may not comply with European Union environmental quality standards.

Wool treated with this product may contain detectable residues: adequate treatment records should be kept and made available, if requested by wool buyers.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin.

When opening the container and using the product, wear elbow-length rubber gloves. Wash hands after use.

FIRST AID INSTRUCTIONS

First aid is not generally required. If in doubt, contact a Poisons Information Centre. Phone Australia 13 11 26.

ADDITIONAL USER SAFETY

Sheep Rehandling Interval: After treatment wait until sheep are dry before rehandling

ENVIRONMENTAL PROTECTION

DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (room temperature). Protect from frost. Store in original container tightly closed in a dry, cool place. DO NOT store in direct sunlight. DO NOT re-use the container. KEEP OUT OF REACH OF CHILDREN.

SAFETY DATA SHEET

For further information please refer to the Safety Data Sheet. For a copy visit our website at www.4farmers.com.au

CONDITIONS OF SALE

4Farmers Australia Pty Ltd and/or its affiliates ("4Farmers") shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on 4Farmer's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely No representative of 4Farmers Australia Pty Ltd has any authority to add to or alter these conditions.

HAZARDS

Avoid breathing fumes, mists, vapours or spray. Do not get in eyes, on skin, or on clothing. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

Hazardous to the Aquatic Environment – Acute Hazard.

Harmful to aquatic life. Avoid release to the environment. Not combustible. Use extinguishing media suited to burning materials. Collect spillage. Dispose of contents and containers as specified on the registered label.

In a Transport Emergency
Dial 000
Police or Fire Brigade
Special Poisons Advice:
13 11 26



OFFICE USE ONLY

Batch Number:

Date of Manufacture:

DOSAGE & ADMINISTRATION

SHAKE WELL BEFORE USE

FOR THE PREVENTION OF BODY STRIKE (Diagram 1)

| SHEEP WEIGHT (KG) | TOTAL VOLUME/SHEEP (ML) | VOLUME PER TREATMENT BAND (ML) | NO. OF BANDS |
|-------------------|-------------------------|--------------------------------|--------------|
| 10-20 | 14 | 7 | 2 |
| 21-30 | 16 | 8 | 2 |
| 31-50 | 20 | 10 | 2 |
| >50 | 24 | 12 | 2 |

FOR THE PREVENTION OF BREECH STRIKE (Diagram 1)

| SHEEP WEIGHT (KG) | TOTAL VOLUME/SHEEP (ML) | VOLUME PER TREATMENT BAND (ML) | NO. OF BANDS |
|-------------------|-------------------------|--------------------------------|--------------|
| 10-20 | 7 | 7 | 1 |
| 21-30 | 8 | 8 | 1 |
| 31-50 | 10 | 10 | 1 |
| >50 | 12 | 12 | 1 |

FOR THE PREVENTION OF FLY STRIKE AFTER MULESING (Diagram 2)

| SHEEP WEIGHT (KG) | TOTAL VOLUME/SHEEP (ML) | VOLUME PER TREATMENT BAND (ML) | NO. OF BANDS |
|-------------------|-------------------------|--------------------------------|--------------|
| Up to 30 | 8 | 4 | 2 |
| 31-50 | 10 | 5 | 2 |

- Apply two bands, one over each side of the mulesing wound and overlapping 25mm onto the surrounding wool.

FOR THE PREVENTION OF FLY STRIKE AFTER MARKING (Diagram 3)

| SHEEP WEIGHT (<30KG) | TOTAL VOLUME/SHEEP (ML) | VOLUME PER TREATMENT BAND (ML) | NO. OF BANDS |
|----------------------|-------------------------|--------------------------------|--------------|
| Ram lambs | 8 | 4 | 2 |
| Ewe lambs | 4 | 4 | 1 |

- When a tail has been amputated, apply 4 ml over the crutch ensuring sufficient coverage of the open wound.
- When the tail has not been amputated, ensure sufficient coverage of the tail where the rubber ring has been applied.

In the case of ram lambs a further 4 ml can be sprayed over and around the scrotum area after the rubber ring has been applied.

FOR THE PREVENTION OF POLL STRIKE (Diagram 1)

- Rams and sheep susceptible to poll strike should be treated. Apply 5 ml of Spray-On to the areas requiring protection, around the poll and horns.

CRITICAL COMMENTS

- Use only a spray-on application gun.
- Ensure the applicator gun is clean.
- Calibrate applicator gun using an accurate volume measure.
- Ensure all areas to be protected are treated.
- When preventing body strike ensure bands overlap along the mid-line.
- When protecting from mulesing wounds from fly strike, take special care to treat the wound and the surrounding 25 mm wool.
- When protecting marking wounds from fly strike, take special care to treat the wound and surrounding wool.
- This product is not recommended for treating existing strikes. Use a product registered as a dressing for existing strikes.
- Heavy rain following the application of this product could diminish the period of protection. DO NOT apply during gusty or windy weather as spray drift may reduce the period of protection.
- 4Farmers Dicyclanil 50g/L Spray-On contains a scourable coloring agent for easy band identification.
- Withholding periods must be observed following application for any purpose, including after mulesing/markings as finite residues will occur.
- It is good agricultural practice for mulesing/markings to be carried out at times of minimal fly activity. Reduced efficacy may occur in sheep affected with dermatophilosis.

Diagram 1






Diagram 2




Diagram 3

